

Macrofauna inhabiting the sponge *Paraleucilla magna* (Porifera: Calcarea) in Rio de Janeiro, Brazil

ANDRÉ PADUA¹, EMILIO LANNA^{1,2} AND MICHELLE KLAUTAU¹

¹Universidade Federal do Rio de Janeiro, Instituto de Biologia, Departamento de Zoologia, Avenida Carlos Chagas Filho, 373, CEP 21941–902, Rio de Janeiro, RJ, Brazil, ²Universidade Federal da Bahia, Instituto de Biologia, Departamento de Biologia Geral, Rua Barão de Jeremoabo, s/n, 40170-175, Campus de Ondina, Salvador, BA, Brasil

Sponges (phylum Porifera) are important components of the benthic marine fauna known for their interactions with vertebrates and a large number of invertebrates seeking for food, shelter or substrate for attachment. Studies on this subject, however, were restricted only to the macrofauna inhabiting sponges of the class Demospongiae. In the present work, we describe the macrofauna associated with a calcareous sponge in Brazil, Paraleucilla magna. Individuals of this allegedly non-native species were monthly collected during one year in Rio de Janeiro (Brazil). Fifty-one taxa representing ten animal phyla were found associated with P. magna. The most frequent and abundant taxa were Crustacea, Mollusca, Polychaeta and Bryozoa, while echinoderms, cnidarians, ascidians, nemerteans, platyhelminthes and sponges were less frequent or even rare and less abundant. Juveniles of several taxa and pregnant females of Crustacea were found associated with P. magna, but these associations were not exclusive. The macrofauna associated with P. magna did not present a clear seasonality, although it was possible to observe a change in the community composition alongside the year. The volume of the sponges was significantly related to the diversity index (H') and number of taxa, but not with evenness (J') and number of individuals. Our results show that P. magna is used as a substrate for attachment and/or shelter by its associates and that most of these associations are just opportunistic. The data presented here reiterate a previous proposal that sponges are important biodiversity reservoirs and that they should be seriously considered in conservation programmes.

Keywords: association, ecology, Crustacea, Mollusca, Polychaeta, Bryozoa

Submitted 23 October 2012; accepted 16 November 2012; first published online 26 February 2013

INTRODUCTION

Sponges (phylum Porifera) have long been considered 'living hotels' due to the great diversity and abundance of other taxonomic groups that are often found in association with them (Pearse, 1950; Klitgaard, 1995; Ribeiro *et al.*, 2003). These associations represent a wide range of ecological interactions, facultative or obligatory, that range from mutualism to parasitism; however, the exact nature of many associations remains unclear (Wulff, 2006). Because sponges have bodies composed of an intricate network of canals, associated organisms may find substrate and shelter inside them (Çinar *et al.*, 2002; Huang *et al.*, 2008). As sponges are important components of benthic communities and interact with a wide range of organisms (Wulff, 2006; Becerro, 2008), they are considered to be important reservoirs of marine biodiversity (Cerrano *et al.*, 2006).

Previous studies of sponge-associated fauna have been carried out in the North Atlantic Ocean (Frith, 1976; Biernbaum, 1981; Peattie & Hoare, 1981; Klitgaard, 1995; Huang *et al.*, 2008; Fiore & Jutte, 2010), the Caribbean (Pearse, 1950; Villamizar &

Laughlin, 1991), the Mediterranean (Rützler, 1976; Koukouras *et al.*, 1985, 1992, 1996; Ilan *et al.*, 1994; Çinar *et al.*, 2002), the Pacific Ocean (Long, 1968; Magnino *et al.*, 1999; Beaulieu, 2001; Skilleter *et al.*, 2005; Cerrano *et al.*, 2006), and the Indian Ocean (Abdo, 2007). Only four studies have been performed in the South Atlantic Ocean: one in Argentina (Cuartas & Excoffon, 1993) and three in Brazil (Duarte & Nalesso, 1996; Ribeiro *et al.*, 2003; Stofel *et al.*, 2008). Other studies along the Brazilian coast have described associations between sponges and particular groups of organisms: parasitic crustaceans (Duarte & Morgado, 1983), decapods (Bezerra & Coelho, 2006), gammarids and caprellids (Serejo, 1998), copepods (Johnsson, 1998, 2000, 2002; Bispo *et al.*, 2006), and polychaetes (Neves & Omena, 2003).

With the exception of two studies that included hexactinellid sponges (Beaulieu, 2001; Fiore & Jutte, 2010), almost all studies of sponge-associated fauna focused on the class Demospongiae. Only one study, conducted in Hampshire, UK, has investigated the associated fauna of a calcareous sponge (Frith, 1976). This study, however, found no fauna associated with either *Sycon ciliatum* (Fabricius, 1780) or *Grantia compressa* (Fabricius, 1780) and did not describe any organisms found with *Leucosolenia botryoides* (Ellis & Solander, 1786) (Frith, 1976).

Paraleucilla magna Klautau *et al.*, 2004 is a calcareous sponge found along the Brazilian coast (adjacent to the

Corresponding author:
Michelle Klautau
Email: mklautau@biologia.ufjf.br




Fig. 1. (A) Location and aerial view of the study area. Vermelha Beach is located at the entrance of the eutrophic Guanabara Bay (GB) (black dot at inferior left corner) (map source: DIVA-GIS, Vermelha Beach photograph: F. Azevedo); (B) *in vivo* photograph of *Paraleucilla magna*.

Rio de Janeiro, São Paulo and Santa Catarina states) and in the Mediterranean (along the southern coast of Italy and around Malta). In both regions, it is considered to be a non-native species, although its origin is still unknown (Klautau *et al.*, 2004; Longo *et al.*, 2007; Zammit *et al.*, 2009; Gravili *et al.*, 2010). It lives attached to hard substrates in photophilous or sciaphilous conditions and in pristine or polluted waters (Klautau *et al.*, 2004; Longo *et al.*, 2007; Gravili *et al.*, 2010). This species has a leuconoid aquiferous system with a large atrial cavity and many canals that can be easily occupied by other organisms. In the original description of *P. magna*, crustaceans, echinoderms, and polychaetes were described as associating with this species (Klautau *et al.*, 2004); however, there has been no subsequent research on its associated fauna. Therefore, to gain knowledge about the associated macrofauna of calcareous sponges, we investigated the composition of macrofauna inhabiting *P. magna* over the course of one year. The objectives of this study were to: (1) describe the species composition of the associated macrofauna of *P. magna* in Rio de Janeiro, south-western Atlantic; (2) investigate the influence

of sponge volume on these associations; and (3) analyse possible seasonal variations of these associations.

MATERIALS AND METHODS

Sampling

Five specimens of *P. magna* (Figure 1B) were collected monthly throughout 2005 (except in February, when only four individuals were collected, and in April, when no collection occurred), totalling 54 specimens. All specimens were collected at Vermelha Beach ($22^{\circ}57'18''\text{ S}-43^{\circ}09'42''\text{ W}$), in Rio de Janeiro, Brazil (Figure 1A; Lanna *et al.*, 2007). Specimens were collected by snorkelling at 0–4 m depth and were removed from the substrate with a knife. While underwater, each specimen was bagged individually (to avoid the escape of associated organisms) and then fixed and preserved in 93% ethanol. At the laboratory, the volume of each sponge was calculated by liquid displacement in a graduated cylinder (see Ribeiro *et al.*, 2003; Lanna *et al.*, 2007). Sponge specimens were then carefully fragmented under a stereomicroscope to remove the macrofauna ($>1\text{ mm}$) that remained inside. Associated organisms of each sponge specimen were separated by morphotype within higher taxa and then identified to the lowest possible taxonomic level with the help of specialists.

Data analysis

We counted the total number of associated individuals and the total number of taxa to calculate species richness, frequency, abundance, density, diversity (H'), and Pielou's evenness (J') (Ludwig & Reynolds, 1988). To investigate whether the total volume of *P. magna* specimens collected each month (i.e. the sum volume of the five analysed individuals) could predict species richness, abundance, diversity and evenness, we performed linear regressions (Sokal & Rohlf, 1995). The values of species richness, frequency, abundance, density, diversity (H'), and Pielou's evenness (J') obtained for each month were used as replicates to test whether these attributes of the associated fauna varied between the dry (April to September) and rainy (October to March) seasons. The rainy season in Rio de Janeiro usually starts in September (Dereczynski *et al.*, 2009), however, we have considered it as starting in October for our analyses because in 2005 (when the specimens were collected) the rainy period started only in that month (AlertaRio, 2011). All data were tested for normality and homoscedasticity prior to performing analyses of variance (ANOVAs). Temporal patterns in the community of associated fauna were assessed by means of a principal component analysis (PCA), in which the dimensionality of 21 species (the number of species that occurred in more than one month) was reduced to only two components (latent variables) representing the primary temporal patterns of dominant species. As most species were rare, and because many zeros were present in the data set (see Table 1), we applied a Hellinger transformation prior to analysis (see Legendre & Gallagher, 2001). PCA scores obtained for each month were used as replicates for the ANOVA to test whether these attributes of the associated fauna varied between the dry and rainy seasons (Jassby & Powell, 1990).

Table 1. Variation of the number of taxa associated with *Paraleucilla magna*. Colonies of Hydrozoa were not quantified, and their presence is marked with 'P'. Total number of individuals/colonies for each taxon and for each month and the total number of taxa of each phylum (within parentheses) are provided. (Por – Porifera, Cni – Cnidaria, Pla – Platyhelminthes, Nem – Nematoda, Ann – Annelida, Art – Arthropoda, Mol – Mollusca, Bry – Bryozoa, Ech – Echinodermata, Asc – Ascidiacea). (*) indicates presence of juveniles.

Taxa	January	February	March	May	June	July	August	September	October	November	December	Total
Por (1) <i>Leucosolenia</i> sp.	1	0	0	0	0	0	0	0	0	0	0	1
Cni (1) Hydrozoa	0	P	0	0	P	0	P	0	0	P	0	
Pla (1) <i>Enchiridium evelinae</i> Marcus, 1949	1	0	0	0	0	0	0	0	0	0	0	1
Nem (1) Unidentified nemertinean	1	0	0	0	0	0	0	0	0	0	0	1
Ann (11) Sabellidae sp.*	1	0	0	0	0	0	0	0	0	0	0	1
<i>Branchiomma luctuosum</i> (Grube, 1869)	1	0	0	0	0	0	0	0	0	0	0	1
Phyllodoceidae	1	0	0	0	0	0	0	0	0	0	0	1
Syllidae sp. 1	3	0	0	0	0	0	0	0	0	0	0	3
Syllidae sp. 2	2	0	0	0	0	0	0	0	1	0	0	3
Syllidae sp. 3	1	0	1	1	1	0	0	0	0	0	0	4
Syllidae sp. 4	0	1	0	0	0	0	0	0	0	0	0	1
Syllidae sp. 5	0	0	0	0	1	0	0	0	0	0	0	1
Syllidae sp. 6	0	0	0	0	0	1	0	0	0	0	0	1
Syllidae sp. 7	0	0	0	0	0	0	1	0	0	0	0	1
<i>Naineris setosa</i> (Verrill, 1900)	0	0	0	0	0	0	0	0	0	0	1	1
Art (17) Stenothoidae	1	0	0	2	0	16	6	7	3	21	2	58
Melitidae sp.	0	0	0	0	0	0	2	0	0	1	0	3
<i>Quadrimaera quadrimana</i> (Dana, 1852)	1	0	0	0	0	0	0	1	0	6	0	8
<i>Elasmopus pecteniscrus</i> (Bate, 1862)	6	0	1	2	1	2	0	0	0	0	3	15
<i>Dulichella appendiculata</i> (Say, 1818)	3	0	0	0	0	0	0	0	0	0	0	3
Podoceridae sp.	0	0	0	0	0	0	0	1	0	0	5	6
Corophiidae	0	0	0	0	0	0	0	0	0	1	0	1
<i>Cymadusa filosa</i> Savigny, 1816	3	0	0	2	0	4	10	3	10	18	1	51
Isopoda	0	0	0	0	0	0	0	0	1	0	0	1
<i>Carpas</i> sp.	0	0	0	0	0	0	0	0	0	3	0	3
<i>Mithrax</i> sp.*	0	0	0	0	0	0	2	0	0	0	0	2
<i>Epialtus bituberculatus</i> Milne Edwards, 1834	0	0	0	0	0	0	1	0	0	0	0	1
<i>Micropanope nuttingi</i> (Rathbun, 1898)	0	0	0	1	0	0	1	0	0	0	0	2
<i>Petrolisthes armatus</i> (Gibbes, 1850)	0	0	0	0	0	0	0	0	0	1	0	1
<i>Pachycheles monilifer</i> (Dana, 1852)	0	0	0	0	0	0	0	0	0	0	2	2
<i>Pachycheles laevidactylus</i> Ortmann, 1892	0	0	0	0	2	5	10	9	4	3	2	35
Pycnogonida sp.	2	0	0	0	0	0	0	0	0	0	0	2
Mol (9) Bivalvia sp.1	0	0	0	1	25	9	3	1	2	1	0	42
Bivalvia sp.2	0	0	0	1	0	0	0	0	0	0	0	1
Bivalvia sp.3	0	0	0	0	1	3	0	1	1	1	2	9
Mytilidae sp.*	0	0	0	3	1	2	0	1	1	2	0	10
<i>Sphenia fragilis</i> (H. Adams & A. Adams, 1854)	0	0	0	1	0	0	0	0	0	0	0	1
<i>Arca</i> sp.	0	0	0	0	0	0	0	0	0	1	0	1
Gastropoda sp.	0	0	0	0	0	0	0	0	0	1	0	1
Calyptraeidae	1	0	1	0	0	1	0	1	0	3	0	7
<i>Crepidula</i> sp.	0	0	1	1	0	0	0	0	0	0	0	2
Bry (3) <i>Bugula neritina</i> (Linnaeus, 1758)	1	0	2	3	4	0	0	0	0	0	0	10
<i>Scrupocellaria</i> aff. <i>reptans</i> (Linnaeus, 1758)	0	1	1	7	5	2	1	0	0	1	2	20
<i>Hippoporina</i> sp.	0	0	1	0	0	0	0	0	0	0	0	1
Ech (4) <i>Ophiactis savignyi</i> (Müller & Troschel, 1842)	1	0	0	0	0	0	1	0	0	0	1	3
<i>Ophiactis lymani</i> Ljungman, 1872	6	0	0	0	0	0	0	0	0	0	3	9
<i>Amphipholis squamata</i> (Delle Chiaje, 1828)	0	0	0	0	0	0	0	0	0	1	0	1
<i>Lytechinus variegatus</i> * (Lamarck, 1816)	0	0	0	0	0	0	0	1	0	0	0	1
Asc (3) <i>Didemnum</i> sp.1	1	1	2	1	1	0	0	0	0	1	1	8
<i>Didemnum</i> sp.2	0	0	2	2	0	1	0	0	0	0	0	5
<i>Botrylloides giganteum</i> (Pérès, 1949)	0	0	0	0	0	1	0	0	0	0	1	2
Number of specimens	38	3	12	28	42	47	38	26	23	66	26	349

Table 2. Summary of the ecological data collected each month.

	January	February	March	May	June	July	August	September	October	November	December	Year
Number of species	20	4	9	14	11	12	12	10	8	18	13	51
Number of specimens	38	3	12	28	42	47	38	26	23	66	26	349
Species diversity (H')	2.7	1.4	2.1	2.4	1.5	2.0	2.1	1.8	1.7	2.1	2.4	3.0
Evenness (J')	0.9	1.0	1.0	0.9	0.6	0.8	0.8	0.8	0.8	0.7	0.9	0.4
Total volume of sponges (cm ³)	115	18	8	11	15	38	52	24	12	21	67	381
Density (ind.cm ⁻³)	0.3	0.2	1.5	2.5	2.8	1.1	0.7	1.1	1.9	3.1	0.4	0.9

RESULTS

Associated macrofauna

A total of 349 individuals, representing 51 species and 10 phyla, were identified living in association with the 54 analysed specimens of *P. magna* (Table 1). The mean species richness of associated taxa was 11.9 species/month (± 4.4 ; Table 2). Arthropoda (mostly Crustacea) showed the highest species richness (17 species); followed by Annelida, with 11 taxa of polychaetes; and Mollusca, with nine species (Table 1). The species diversity of the total associated macrofauna was high ($H' = 3.0$), but the total evenness was low ($J' = 0.4$) (Table 2).

The most abundant higher taxa were Arthropoda (54%), Mollusca (21%), and Bryozoa (9%) (Figure 2), while the most frequent were Arthropoda, Annelida (Polychaeta), Mollusca, and Bryozoa, present in 72.2%, 57.4%, 48.2%, and 40.7% of the sponges, respectively (Figure 3). Chordata (Ascidiacea), Cnidaria (Hydrozoa), and Echinodermata were found less frequently (present in 22.2%, 14.8%, and 12.9% of the sponges, respectively), while Platyhelminthes, Nemertea, and Porifera were found in only 1.8% of specimens (Figure 3). The density of associated individuals was highest in November and June (3.1 and 2.8 ind.cm⁻³) and lowest in February and January (0.2 and 0.3 ind.cm⁻³). This variation was not significantly different between the dry and rainy seasons (Table 3A).

Juvenile representatives of Crustacea (*Mithrax* sp.), Polychaeta (*Sabellidae* sp., *Syllidae* spp.), Mollusca (*Mytilidae* sp.), and Echinodermata (*Lytechinus variegatus*)




Fig. 2. Proportion of the higher taxa associated with *Paraleucilla magna*.




Fig. 3. Frequency (%) of the phyla associated with *Paraleucilla magna*.

were found living associated with *P. magna*. In addition, pregnant crustacean females were also frequently observed.

Volume

Total sponge volume (i.e. the sum volume of sponges collected each month; Table 2) did not differ between seasons

Table 3. Summary of the ANOVA testing the influence of seasonality (dry versus rainy seasons) on community descriptors during the study period ($df =$ degrees of freedom; Sum Sq = sum of squares; Mean Sq = mean of squares; $Pr > F = P$ value associated with the F statistic; * $P < 0.05$).

	df	Sum Sq	Mean Sq	F value	Pr > F
(A) Density of associated individuals					
Season	1	0.552	0.5520	0.4402	0.5236
Residuals	9	11.285	1.2539		
(B) Sponge volume					
Season	1	404.600	404.6000	0.3559	0.5655
Residuals	9	10232.800	1137.0000		
(C) Species richness					
Season	1	0.109	0.1091	0.0049	0.9455
Residuals	9	198.800	22.0889		
(D) Number of associated individuals					
Season	1	186.380	186.3800	0.5943	0.4605
Residuals	9	2822.530	313.6200		
(E) Diversity (H')					
Season	1	0.032	0.03292	0.1850	0.6773
Residuals	9	1.601	0.17797		
(F) Evenness					
Season	1	0.025	0.02579	2.6065	0.1409
Residuals	9	0.089	0.00989		




Fig. 4. Monthly variation of the number of species and individuals associated with *Paraleucilla magna*.

(Table 3B) but was significantly correlated to both species diversity (H') ($R^2 = 0.43$, $df = 10$, $P = 0.027$, Figure 5A) and the number of taxa (species richness) ($R^2 = 0.37$, $df = 10$, $P = 0.04$, Figure 5B), indicating that larger sponges contained a higher variety of taxa and a higher diversity of species. Nonetheless, regression analyses indicated that the total volume in each month did not correlate to either the Pielou evenness index (J') ($R^2 = 0.04$, $df = 10$, $P = 0.52$, Figure 5C) or the total number of associated individuals (abundance) ($R^2 = 0.03$, $df = 10$, $P = 0.56$, Figure 5D).

Seasonality

The periods of lowest and highest species richness (February = 4; January = 20, respectively) coincided with the months of lowest and highest diversity (H') (February - $H' = 1.4$; January - $H' = 2.7$) (Table 2). Abundance (i.e. the number of associated individuals) was lowest in February (only three individuals), and highest in November (66 individuals) (Table 1; Figure 4). The evenness of associated

macrofauna tended to be high, being highest in February and March ($J' = 1.0$) and lowest in June ($J' = 0.6$) (Table 2). None of these community descriptors differed significantly between the dry and rainy seasons (Table 3C–F).

Seasonal changes in the community of macrofauna associated with *P. magna* were analysed using biplots based on PCA (Figure 6A). The total amount of variation explained by the first two scores (corresponding to the first two principal components) was 56.9%. The PCA biplot did not show a clear seasonal difference between the dry and rainy seasons. Nevertheless, three groups of species were partially distinguished by the analysis:

Group A (formed mainly by the bryozoan *Scrupocellaria* aff. *reptans* (Linnaeus, 1758) and the ascidians *Didemnum* sp. 1 and *Bugula neritina* (Linnaeus, 1758)), which appeared between February and June;

Group B (formed mainly by the mollusc *Bivalvia* sp.1 and the crustaceans *Pachycheles laevidactylus* Ortmann, 1892 and *Cymadusa filosa* Savigny, 1816) that appeared from July to November;

Group C (formed mainly by the ophiuroid *Ophiactis lymani* Ljungman, 1872) comprised only one species and was found exclusively in January and December.

Scores of the first component (PC1), which account for 38.7% of the variation, did not differ significantly between the dry and rainy seasons (Figure 6B; Table 4A). However, the scores of the second component (PC2), which account for 18.2% of the variation, were significantly different between these seasons (Figure 6C; Table 4B).

DISCUSSION

Paraleucilla magna exhibited moderate-to-low richness of associated macrofauna (51 species) relative to all other




Fig. 5. Quantitative analyses of the macrofauna associated with *Paraleucilla magna*. Linear regression between sponge volume and (A) species diversity (H'); (B) number of taxa; (C) evenness (J'); (D) number of individuals. The dotted lines indicate the 95% confidence intervals.




Fig. 6. Principal component analysis (PCA) of the associated fauna of *Paraleucilla magna*: (A) biplot representation of the PCA showing both observations (months) and variables (species) in the same graph. The left and bottom axes use the unity for observations, while the top and right axes are graduated according to the first two principal components of the original variables. PC1 accounts for 38.7% of the total variation, while PC2 accounts for 18.2%. Months are represented by upper case letters (Species: bi1, Bivalvia sp. 1; bi3, Bivalvia sp. 3; bot, *Botrylloides giganteum*; bug, *Bugula neritina*; cal, Calyptraeidae; cre, *Crepidula* sp.; cym, *Cymadusa filosa*; di1, *Didemnum* sp. 1; di2, *Didemnum* sp. 2; ela, *Elasmopus pecteniscrus*; mel, Melitidae sp.; mic, *Micropanope nuttingi*; myt, Mytilidae sp.; opl, *Ophiactis lymani*; ops, *Ophiactis savignyi*; pac, *Pachycheles laevidactylus*; pod, Podceridae sp.; qua, *Quadrimaera quadrimana*; scr, *Scrupocellaria aff. reptans*; sy2, Syllidae sp. 2; sy3, Syllidae sp.); (B and C) box plots of the scores of (B) the first principal component (PC1) and (C) the second principal component (PC2) in the dry and rainy seasons. Each box displays the median, upper and lower quartiles of the distribution of sponge volume per month. Box whiskers represent the maximum and minimum range, while empty circles show outliers.

sponge species investigated to date (48 Demospongiae and two Hexactinellida). Demosponges, for example, yielded an average of 95.5 associated taxa per species (± 162.2 of standard

deviation), with a minimum of two and a maximum of 809 taxa (e.g. Westinga & Hoetjes, 1981; Villamizar & Laughlin, 1991; Cuartas & Excoffon, 1993; Klitgaard, 1995; Koukouras *et al.*, 1996; Betancourt-Lozano *et al.*, 1998; Magnino *et al.*, 1999; Çinar *et al.*, 2002; Neves & Omena, 2003; Ribeiro *et al.*, 2003; Abdo, 2007; Huang *et al.*, 2008). In *P. magna*, Crustacea was the most abundantly represented group of associated organisms (54%), followed by Mollusca (21%), and Bryozoa (9%). In other studied sponges, Crustacea was also one of the two most abundantly represented groups, being present in 80% of the sponge species examined, followed by Polychaeta (60%) and Echinodermata (24%). Molluscs were the second most abundant group in *P. magna* (21%); however, this is not a common occurrence, as they have been identified as a dominant group in only a few species of sponges (8% of those examined so far; Long, 1968; Peattie & Hoare, 1981; Klitgaard, 1995; Koukouras *et al.*, 1996). The same pattern

Table 4. Summary results of the analysis of variance carried out on the scores of the two main principal components testing the seasonality (dry versus rainy seasons) of the associated fauna during the studied period (df = degrees of freedom; Sum Sq = sum of squares; Mean Sq = mean of squares; Pr > F = P value associated with the F statistic; *P < 0.05).

	df	Sum Sq	Mean Sq	F value	Pr > F
(A) PC1					
Season	1	0.2126	0.2126	0.9401	0.3576
Residuals	9	2.0353	0.2261		
(B) PC2					
Season	1	0.3972	0.3972	5.4382	0.0446*
Residuals	9	0.6573	0.0730		

occurs with Bryozoa, which was the third most abundant taxon in *P. magna* but is not considered to be among the two most abundant organisms in other studied sponges. However, bryozoans were the second most dominant group (12.8% of the total number of taxa) found in demosponges of the Faroe Island, north-eastern Atlantic (Klitgaard, 1995) and, as in the present study, Klitgaard (1995) also found that most of the bryozoans were attached to the outer surface of the sponges. Associations between sponges and bryozoans may be related to the fact that sponges may provide suitable substrate to bryozoans in habitats of otherwise limited substrate availability, as noted by Klitgaard (1995).

A study of the associated fauna of the demosponge *Mycale microsigmatosa* Arndt, 1927 was performed at the same location of the present study (Ribeiro *et al.*, 2003). Both *P. magna* and *M. microsigmatosa* exhibit associated macrofauna of similar species richness (51 and 75 species, respectively) and composition. However, the differences observed in taxonomic composition between these two sympatric species can be explained by the different sample sizes of each study. In the present study, we analysed 54 specimens of *P. magna*, while Ribeiro *et al.* (2003) analysed 19 specimens of *M. microsigmatosa*. Species diversity was the same in *P. magna* and *M. microsigmatosa* ($H' = 3.0$), while evenness was lower in *P. magna* ($J' = 0.4$, versus $J' = 0.7$ for *M. microsigmatosa*). The difference in evenness values between both species may be also due to sampling differences. In the present work, several collections throughout the year were made, while Ribeiro *et al.* (2003) made only one collection. The most striking difference between these two species is in the total number of associated individuals (abundance): *P. magna* was associated with 349 individuals (0.9 ind.cm^{-3}), while *M. microsigmatosa* was associated with 2235 (13 ind.cm^{-3}). If we consider that both sponges have the same type of aquiferous system (leuconoid), we could expect similar internal canals and, consequently, similar associated macrofauna. Nonetheless, *P. magna* has a large atrium, while *M. microsigmatosa* has only canals, and whereas *P. magna* is massive, *M. microsigmatosa* is an incrustant sponge. In addition, the external surface of *P. magna* is full of folds, while *M. microsigmatosa* has a smoother surface. Despite these morphological characteristics that seem to characterize *P. magna* as a better host, *M. microsigmatosa* is

host to more associated organisms. A possible explanation for this difference in macrofauna abundance is the presence of chemicals that might reduce predation in *M. microsigmatosa* and, consequently, provide more protection for its associated macrofauna. Although this hypothesis has not been tested, *M. microsigmatosa* does produce a series of compounds, some of which inhibit microorganism proliferation (Compagnone *et al.*, 1999; Marinho *et al.*, 2009, 2010; Santos *et al.*, 2010). The potential importance of sponge allelochemicals in influencing the composition and abundance of associated fauna has already been pointed out (Koukouras *et al.*, 1992; Skilleter *et al.*, 2005). A good example can be found in the work of Betancourt-Lozano *et al.* (1998), which describes a significant relationship between inquilinism and the antibiosis activity of *Aplysina fistularis* (Pallas, 1766) in Mexico.

Paraleucilla magna shares with *M. microsigmatosa* at least three associated species, two of which (the ophiuroids *Amphipholis squamata* and *Ophiactis savignyi*) occur commonly in other sponge species (Table 5). Although echinoderms have been found in only 12.9% of the analysed specimens of *P. magna*, they (particularly Ophiuroidea) are commonly found in demosponges (Wendt *et al.*, 1985; Duarte & Nalesso, 1996; Betancourt-Lozano *et al.*, 1998; Ribeiro *et al.*, 2003; Clavico *et al.*, 2006; Abdo, 2007) and other benthic organisms, such as bryozoans (Morgado & Tanaka, 2001). Associations of *Ophiactis savignyi* and *O. lymani* with marine organisms are apparently common. For example, both species have been described as common epifauna on the tubes of the polychaete *Phyllochaetopterus socialis* Claparède, 1869 (Nalesso *et al.*, 1995), on the octocoral *Carijoa riisei* (Duchassaing & Michelotti, 1860) (Neves *et al.*, 2007), and on algae (Mladenov & Emson, 1988). The frequent association of these ophiuroid species with varied taxa (algae, polychaetes, corals and sponges) may indicate that these associations (including with *P. magna*) are only occasional or opportunistic. These ophiuroids may seek out these organisms only for protection or food (Klitgaard, 1995).

The volume of *P. magna* was positively related only to species diversity and number of taxa (richness). These relationships have already been observed in other sponge species: *S. foetidus* (for species diversity) and *M. microsigmatosa*, *M. angulosa*, *S. foetidus*, and *Spheciospongia vesparium*

Table 5. Species associated with *Paraleucilla magna* that were already found associated with other sponge species. 1 – *Mycale microsigmatosa* (Rio de Janeiro, Brazil; Ribeiro *et al.*, 2003); 2 – *Mycale angulosa* (São Paulo, Brazil; Duarte & Nalesso, 1996); 3 – *Dysidea robusta* Vilanova & Muricy, 2001 (Rio de Janeiro, Brazil; Serejo, 1998); 4 – *Topsentia* sp. (south-eastern United States; Fiore & Jutte, 2010); 5 – *Ircinia campana* (Lamarck, 1814) (south-eastern United States; Fiore & Jutte, 2010); 6 – *Sarcotragus foetidus* (Turkish Aegean coast; Çinar *et al.*, 2002); 7 – *Aplysina lacunosa* (Pallas, 1766) (Venezuelan Caribbean; Villamizar & Laughlin, 1991); 8 – *Sarcotragus fasciculatus* (Pallas, 1766) (North Aegean Sea; Koukouras *et al.*, 1985); 9 – *Sidonops corticosylifera* (Hajdu, Muricy, Custodio, Russo & Peixinho, 1992) (Rio de Janeiro, Brazil; Clavico *et al.*, 2006); 10 – *Halichondria panicea* (Pallas, 1766) (Menai Strait, UK; Peattie & Hoare, 1981); 11 – *Ircinia strobilina* (Lamarck, 1816) (Bimini, Bahamas; Pearse, 1950); 12 – *Geodia macandrewii* Bowerbank, 1858 (Faroe Islands; Klitgaard, 1995); 13 – *Cliona varians* (Duchassaing & Michelotti, 1864) (Stofel *et al.*, 2008).

Associated species	1	2	3	4	5	6	7	8	9	10	11	12	13
Crustacea													
<i>Elasmopus pecteniscus</i>	x												
<i>Quadrimaera quadrimana</i>			x										
<i>Dulichieilla appendiculata</i>			x	x	x								
<i>Cymadusa filosa</i>													x
Echinodermata													
<i>Amphipholis squamata</i>	x	x						x		x		x	
<i>Ophiactis lymani</i>		x											x
<i>Ophiactis savignyi</i>	x	x				x	x		x		x		

(Lamarck, 1815) (for richness) (Westinga & Hoetjes, 1981; Duarte & Nalesso, 1996; Çinar et al., 2002; Ribeiro et al., 2003). In *P. magna*, higher volumes can reflect a diverse array of microhabitats inside the sponge, such as more and larger folds, or larger atria and oscula, which could accommodate larger organisms and, consequently, a higher diversity of taxa. On the other hand, no relationship between volume and number of individuals was observed in *P. magna*, and this relationship has also not been observed in several demosponge species (four from the Aegean Sea, Koukouras et al., 1992; and two from Australia, Skilleter et al., 2005). In *P. magna*, large volumes might provide habitat for other species that could then compete with the fauna that live in smaller sponges. The fact that we found associated organisms in a great variety of sponge volumes (from 0.3 cm³ to 37 cm³) suggests that this species is rapidly colonized by organisms in the environment.

In the present study, no significant seasonal variation in community descriptors of the fauna associated with *P. magna* (species richness, number of individuals, species diversity (H'), and evenness index (J')) was detected. This lack of seasonal variation can be explained, in part, by the relationship of some of these descriptors with sponge volume (as described above). As neither sponge volume nor the community descriptors exhibit variation between the dry and rainy seasons (see Table 2), the absence of any seasonal trend could be expected. However, it is important to consider that sample size, differences in the sponges volume collected each month and a possible atypical year could have influenced these results.

Although the PCA biplot (Figure 6A) suggests no seasonal variation between the dry and rainy seasons, the second component (PC2) scores differed significantly between seasons. This latter result indicates that some environmental change (in features such as salinity, temperature, or food availability) might influence the composition of the associated fauna community. However, the causes of variation explained by the first component (PC1) are unknown and not likely to be correlated with season. On the other hand, we observed three groups of species that occupied *P. magna* in temporal succession (Groups A, B, and C). The establishment of these groups may reflect the life cycle of the associated organisms.

We frequently found pregnant crustacean females and juveniles of several taxa (molluscs, crustaceans, echinoderms, and polychaetes) inhabiting *P. magna* that probably used their host as a temporary shelter during vulnerable periods of their life cycle (i.e. reproductive or juvenile stages). This kind of relationship can be characterized as opportunistic. Ribeiro et al. (2003) and Abdo (2007) also found pregnant females, juveniles or reproductively active individuals associated with *M. microsigmata* and two *Haliclona* species in Brazil and Australia, respectively.

These findings suggest that sponges may be important shelters during some stages of the life cycle of many invertebrates, enhancing their survival. All of these aspects regarding the role of sponges in the community reiterate a previous proposal (Cerrano et al., 2006) namely, that sponges are important reservoirs of biodiversity and that the phylum Porifera should be seriously considered in conservation programs.

ACKNOWLEDGEMENTS

We thank the taxonomy specialists for helping us with identifications of the associated fauna: Luciana Muguet (Bryozoa),

Cléo Oliveira (Mollusca), Juliana Bahia (Platyhelminthes), Carlos Renato R. Ventura and Fernanda Viana (Echinodermata), Daniela Barbosa (Asciadiacea), Paulo Paiva (Polychaeta) and Tereza G. Silva (Crustacea). We are thankful to Baslavi Condor and Paulo Paiva for their help in data analysis and to the anonymous referees for their suggestions that improved manuscript quality. We thank the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), the Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ) (E-26/111.541/2008), and the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq-480368/2008 2; CNPq/PIBIC-480368/2008 2) for grants and fellowships during this project.

REFERENCES

- Abdo D.A. (2007) Endofauna differences between two temperate marine sponges (Demospongiae; Haplosclerida; Chalinidae) from southwest Australia. *Marine Biology* 152, 845–854.
- AlertaRio S. (2011) Sistema AlertaRio, Prefeitura do Rio de Janeiro, RJ – www.rio.rj.gov.br/alertario (accessed 10 August 2011).
- Beaulieu S.E. (2001) Life on glass houses: sponge stalk communities in the deep sea. *Marine Biology* 138, 803–817.
- Becerro M.A. (2008) Quantitative trends in sponge ecology research. *Marine Ecology* 29, 167–177.
- Betancourt-Lozano M., Gonzalez-Farias F., Gonzalez-Acosta B., Garcia-Gasca A. and Bastida-Zavala J.R. (1998) Variation of antimicrobial activity of the sponge *Aplysina fistularis* (Pallas, 1766) and its relation to associated fauna. *Journal of Experimental Marine Biology and Ecology* 223, 1–18.
- Bezerra L.E.A. and Coelho P.A. (2006) Crustáceos decápodos associados a esponjas no litoral do estado do Ceará, Brasil. *Revista Brasileira de Zoologia* 23, 699–702.
- Biernbaum C.K. (1981) Seasonal changes in the amphipod fauna of *Microciona prolifera* (Ellis and Solander) (Porifera, Demospongia) and associated sponges in a shallow salt marsh creek. *Estuaries* 4, 85–96.
- Bispo R., Johnsson R. and Neves E. (2006) A new species of *Astrocheres* (Copepoda, Siphonostomatoida, Astrocheridae) associated to *Placospongia cristata* Boury-Esnault (Porifera) in Bahia State, Brazil. *Zootaxa* 1351, 23–34.
- Cerrano C., Calcinaï B., Pinca S. and Bavestrello G. (2006) Reef sponges as host of biodiversity: cases from North Sulawesi. In Suzuki Y., Nakamori T., Hidaka M., Kayanne H., Casareto B.E., Nadao K., Yamano H. and Tsuchiya M. (eds), *10th International Coral Reef Symposium Proceedings, Okinawa, 28 June–2 July 2006. Functional roles of sponges in coral reefs*. Tokyo: Japanese Coral Reef Society, pp. 208–213.
- Çinar M.E., Katagan T., Ergen Z. and Sezgin M. (2002) Zoobenthos inhabiting *Sarcotragus muscarum* (Porifera: Demospongiae) from the Aegean Sea. *Hydrobiologia* 482, 107–117.
- Clavico E.E.G., Muricy G., da Gama B.A.P., Batista D., Ventura C.R.R. and Pereira R.C. (2006) Ecological roles of natural products from the marine sponge *Geodia corticosylifera*. *Marine Biology* 148, 479–488.
- Compagnone R.S., Oliveri M.C., Piña I.C., Marques S., Rangel H.R., Dagger F., Suárez A.I. and Gómez M. (1999) 5-Alkylpyrrole-2-Carboxaldehydes from the Caribbean sponges *Mycale microsigmata* and *Desmapsamma anchorata*. *Natural Product Letters* 13, 203–211.

- Cuarteras E.I. and Excoffon A.C. (1993) La fauna acompañante de *Hymeniacidon sanguinea* (Grant, 1827) (Porifera: Demospongiae). *Neotrópica* 39, 3–10.
- Dereczynski C.P., Oliveira J.S. and Machado C.O. (2009) Climatologia da precipitação no município do Rio de Janeiro. *Revista Brasileira de Meteorologia* 24, 24–38.
- Duarte L.F.L. and Morgado E.H. (1983) Crustáceos parasitos de invertebrados associados à esponja *Zygomycale parishii* (Bowerbank) e ao briozoário *Schizoporella unicornis* (Jonhston, 1847). *Iheringiasérie Zoologia* 62, 3–11.
- Duarte L.F.L. and Nalesso R. C. (1996) The sponge *Zygomycale parishii* (Bowerbank) and its endobiotic fauna. *Estuarine, Coastal and Shelf Science* 42, 139–151.
- Fiore C.L. and Jutte P.C. (2010) Characterization of macrofaunal assemblages associated with sponges and tunicates collected off the south-eastern United States. *Invertebrate Biology* 129, 105–120.
- Frith D.W. (1976) Animals associated with sponges at North Hayling, Hampshire. *Zoological Journal of the Linnean Society* 58, 353–362.
- Gravili C., Belmontea G., Cecere E., Denitto F., Giangrande A., Guidetti P., Longo C., Mastrototaro F., Moscatello S., Petrocelli A., Piraino S., Terlizzi A. and Boero F. (2010). Nonindigenous species along the Apulian coast, Italy. *Chemistry and Ecology* 26, 121–142.
- Huang J.P., McClintock J.B., Amsler C.D. and Huang Y.M. (2008) Mesofauna associated with the marine sponge *Amphimedon viridis*. Do its physical or chemical attributes provide a prospective refuge from fish predation? *Journal of Experimental Marine Biology and Ecology* 362, 95–100.
- Ilan M., Ben-Eliahu M.N. and Galil B.S. (1994) Three deep water sponges from the eastern Mediterranean and their associated fauna. *Ophelia* 39, 45–54.
- Jassby A.D. and Powell T.M. (1990) Detecting changes in ecological time series. *Ecology* 71, 2044–2052.
- Johnsson R. (1998) Six new species of the genus *Astrocheres* (Copepoda: Siphonostomatoida) associated with sponges in Brazil. *Nauplius* 6, 61–99.
- Johnsson R. (2000) *Spongiopsyllus adventicius* new species and genus of Entomolepididae (Copepoda: Siphonostomatoida) associated with sponges in Brazil. *Hydrobiologia* 417, 115–119.
- Johnsson R. (2002) Astrocherids (Copepoda: Siphonostomatoida) associated with invertebrates from California Reefs: Abrolhos (Brazil). *Hydrobiologia* 470, 247–266.
- Klautau M., Monteiro L. and Borojevic R. (2004) First occurrence of the genus *Paraleucilla* (Calcarea, Porifera) in the Atlantic Ocean: *P. magna* sp. nov. *Zootaxa* 710, 1–8.
- Klitgaard A.B. (1995). The fauna associated with outer shelf and upper slope sponges (Porifera, Demospongiae) at the Faroe Islands, Northeastern Atlantic. *Sarsia* 80, 1–22.
- Koukouras A., Voultziadou-Koukouras E., Chintiroglou H. and Dounas C. (1985) Benthic bionomy of the North Aegean Sea. 3. A comparison of the macrobenthic animal assemblages associated with 7 sponge species. *Cahiers de Biologie Marine* 26, 301–319.
- Koukouras A., Russo A., Voultziadou-Koukouras E., Dounas C. and Chintiroglou C. (1992) Relationship of sponge macrofauna with the morphology of their hosts in the North Aegean Sea. *Internationale Revue der Gesamten Hydrobiologie* 77, 609–619.
- Koukouras A., Russo A., Voultziadou-Koukouras E., Arvanitidis C. and Stefanidou D. (1996) Macrofauna associated with sponge species of different morphology. *Marine Ecology—Pubblicazioni Della Stazione Zoologica Di Napoli* 17, 569–582.
- Lanna E., Monteiro L.C. and Klautau M. (2007) Life cycle of *Paraleucilla magna* Klautau, Monteiro and Borojevic, 2004 (Porifera, Calcarea). In Custódio M.R. Lôbo-Hajdu G. Hajdu E. and Muricy G. (eds) *Porifera research—biodiversity, innovation and sustainability*. Rio de Janeiro: Museu Nacional—Série Livros 28, pp. 413–418.
- Legendre P. and Gallagher E. (2001) Ecologically meaningful transformations for ordination of species data. *Oecologia* 129, 271–280.
- Long E.R. (1968) The associates of four species of marine sponges of Oregon and Washington. *Pacific Science* 22, 347–351.
- Longo C., Mastrototaro F. and Corriero G. (2007) Occurrence of *Paraleucilla magna* (Porifera: Calcarea) in the Mediterranean sea. *Journal of the Marine Biological Association of the United Kingdom* 87, 1749–1755.
- Ludwig J.A. and Reynolds J.F. (1988) *Statistical ecology—a primer on methods and computing*. New York: Wiley-Interscience.
- Magnino G., Pronzato R., Sarà A. and Gaino E. (1999) Fauna associated with the horny sponge *Anomoianthella lamella* Pulitzer-Finali & Pronzato, 1999 (Ianthellidae, Demospongiae) from Papua-New Guinea. *Italian Journal of Zoology* 66, 175–181.
- Marinho P.R., Moreira A.P.B., Pellegrino F.L.P.C., Muricy G., Bastos M.C.F., Santos K.R.N., Giambiagi-deMarval M. and Laport M.S. (2009) Marine *Pseudomonas putida*: a potential source of antimicrobial substances against antibiotic-resistant bacteria. *Memórias do Instituto Oswaldo Cruz, Rio de Janeiro* 104, 678–682.
- Marinho P.R., Muricy G.R.S, Silva M.F.L., deMarval M.G. and Laport M.S. (2010) Antibiotic-resistant bacteria inhibited by extracts and fractions from Brazilian marine sponges. *Brazilian Journal of Pharmacognosy* 20, 267–275.
- Mladenov P.V. and Emson R.H. (1988) Density, size structure and reproductive characteristics of fissiparous brittle stars in algae and sponges: evidence for interpopulational variation in levels of sexual and asexual reproduction. *Marine Ecology Progress Series* 42, 181–194.
- Morgado E.H. and Tanaka M.O. (2001) The macrofauna associated with the bryozoan *Schizoporella errata* (Walters) in southeastern Brazil. *Scientia Marina* 65, 173–181.
- Nalesso R.C., Duarte L.F.L., Pierozzi I. and Enumo E.F. (1995) Tube epifauna of the Polychaete *Phyllochaetopterus socialis* Claparède. *Estuarine, Coastal and Shelf Science* 41, 91–100.
- Neves B.M., Lima E.J.B. and Pérez C.D. (2007) Brittle stars (Echinodermata: Ophiuroidea) associated with the octocoral *Carijoa riisei* (Cnidaria: Anthozoa) from the littoral of Pernambuco, Brazil. *Journal of the Marine Biological Association of the United Kingdom* 87, 1263–1267.
- Neves G. and Omena E. (2003) Influence of sponge morphology on the composition of the polychaete associated fauna from Rocas Atoll, northeast Brazil. *Coral Reefs* 22, 123–129.
- Pearse A. S. (1950) Notes on the inhabitants of certain sponges at Bimini. *Ecology* 31, 149–151.
- Peattie M. E. and Hoare R. (1981) The sublittoral ecology of the Menai Strait. 2. The sponge *Halichondria panicea* (Pallas) and its associated fauna. *Estuarine, Coastal and Shelf Science* 13, 621–635.
- Ribeiro S.M., Omena E.P. and Muricy G. (2003) Macrofauna associated to *Mycale microsigmatosa* (Porifera, Demospongiae) in Rio de Janeiro State, SE Brazil. *Estuarine, Coastal and Shelf Science* 57, 951–959.
- Rützler K. (1976) Ecology of Tunisian commercial sponges. *Tethys* 7, 249–264.
- Santos O.C.S., Pontes P.V.M.L., Santos J.F.M., Muricy G., Giambiagi-deMarval M. and Laport M.S. (2010) Isolation, characterization and phylogeny of sponge-associated bacteria with antimicrobial activities from Brazil. *Research in Microbiology* 161, 604–612.

- Serejo C.S.** (1998) Gammaridean and caprellidean fauna (Crustacea) associated with the sponge *Dysidea fragilis* Johnston at Arraial do Cabo, Rio de Janeiro, Brazil. *Bulletin of Marine Science* 63, 363–385.
- Skilleter G.A., Russell B.D., Degnan B.M. and Garson M.J.** (2005) Living in a potentially toxic environment: comparisons of endofauna in two congeneric sponges from the Great Barrier Reef. *Marine Ecology Progress Series* 304, 67–75.
- Sokal R.R. and Rohlf F.J.** (1995) *Biometry*. San Francisco, CA, Freeman & Co.
- Stofel C.B., Canton G.C., Antunes L.A.S. and Eutrópio F.J.** (2008) Fauna associada à esponja *Cliona varians* (Porifera, Demospongiae). *Natureza Online* 6, 16–18.
- Villamizar E. and Laughlin R.A.** (1991) Fauna associated with the sponges *Aplysina archeri* and *Aplysina lacunosa* in a coral reef of the Archipiélago de Los Roques, National Park, Venezuela. In Reitner J. and Keupp H. (eds) *Fossil and recent sponges*. Berlin: Springer, pp. 522–542.
- Wendt P.H., van Dolah R.F. and O'Rourke C.B.** (1985) A comparative study of the invertebrate macrofauna associated with seven sponge and coral species collected from the South Atlantic Bight. *Journal of the Elisha Mitchell Scientific Society* 101, 187–203.
- Westinga E. and Hoetjes P.C.** (1981) The intrasponge fauna of *Spherospongia vesparia* (Porifera, Demospongiae) at Curaçao and Bonaire. *Marine Biology* 62, 139–150.
- Wulff J.L.** (2006) Ecological interactions of marine sponges. *Canadian Journal of Zoology* 84, 146–166.
- and
- Zammit P.P., Longo C. and Schembri P.J.** (2009) Occurrence of *Paraleucilla magna* Klautau et al., 2004 (Porifera: Calcarea) in Malta. *Mediterranean Marine Science* 10, 135–138.

Correspondence should be addressed to:

Michelle Klautau
Universidade Federal do Rio de Janeiro, Instituto de
Biologia, Departamento de Zoologia, Avenida Carlos Chagas
Filho, 373, CEP 21941–902, Rio de Janeiro, RJ, Brazil
e-mail: mklautau@biologia.ufrj.br