

GEOCHRONOLOGY (U-PB/PB-PB) AND ISOTOPIC SIGNATURE (RB-SR/SM-ND) OF THE PALEOPROTEROZOIC GUANAMBI BATHOLITH, SOUTHWEST BAHIA STATE (NE BRAZIL)

MARIA DE LOURDES DA SILVA ROSA¹, HERBET CONCEICAO¹, FELIX OBERLI², MARTIN MEIER², HERVE MARTIN³, MOACIR JOSE BUENANO MACAMBIRA⁴, EMERSON BARRETO DOS SANTOS¹, MARCIO MATTOS PAIM¹, GERALDO AUGUSTO DA SILVA LEAHY¹ AND LUIZ ROGERIO BASTOS LEAL¹

ABSTRACT The Guanambi batholith (GB) is located in south-western Bahia, Brazil, where it forms part of the Urandi-Paratinga mobile belt. The batholith extends over an area of almost 6,000 km² and is mainly composed of syenites and monzonites, with subordinate granites, mafic syenites and lamprophyric dykes. Two main units can be recognized: the multiple intrusions, being divided into four regions (Paratinga, Laguna, Igapora and Guanambi) and the so-called late-intrusions forming the Cara Suja, Cerafina and Estreito massifs. U-Pb and Pb-Pb isotopic data demonstrate that the batholith was emplaced 2.05 Ga ago. GB has (i) highly negative initial ϵ_{Nd} (-7.4 to -10.6), pointing to an enriched source, and (ii) Sr initial ratios between 0.704 and 0.707. Both chemical and isotopic signatures suggest that GB formed by fractional crystallization of a lamprophyric magma derived from a Paleoproterozoic enriched mantle source.

Keywords: Guanambi batholith, geochronology, enriched mantle, syenites

INTRODUCTION Syenitic magmatism, widespread in Bahia State (Brazil) during Precambrian times, can be divided into two age groups: Transamazonian (1.8-2.1 Ga) and Brasiliano (0.45-0.70 Ga). Ca. 90% of the syenite massifs in Bahia belong to the Transamazonian group. These silica-saturated rocks form large batholiths, which intrude the Paleoproterozoic mobile belts representing part of the basement of the Sao Francisco Craton. <10% of the syenites belong to the Brasiliano group, represented by small ring complexes dominated by plutonic silica-undersaturated rocks.

There are two Paleoproterozoic mobile belts in Bahia: i) the Urandi-Paratinga mobile belt (UPMB) and ii) the Salvador-Curaçá mobile belt (SCMB) located in the western and eastern parts of the state, respectively. Both belts are characterized by the emplacement of alkali-syenitic, potassic to ultrapotassic rocks during late-stage stabilization (Conceição 1993). The syenites of the SCMB are concentrated in four separate massifs, being emplaced as tabular bodies and displaying gneissic texture, whereas in the UPMB, the syenites form a large, 6,000 km² sized, single batholith, the Guanambi batholith (GB), which is related to a pull-apart system and shows exclusively magmatic textures. In this paper, we focus on the geochronology and isotopic signature of the Guanambi Batholith, which represents the largest association of potassic-ultrapotassic syenitic and monzonitic rocks in the São Francisco Craton.

GEOLOGICAL SETTING The north-south trending GB has a sigmoidal shape and is ca. 220 km long and between 15 to 60 km wide (Fig. 1). It is located in the southern part of the Sao Francisco Craton in the Urandi-Paratinga mobile belt (UPMB). The belt consists of three units (Fig. 1): (i) the Santa Izabel Complex at its eastern border, represented by gneisses, migmatites and granulites with U-Pb zircon ages of ca. 3.35 ± 0.29 Ga (Rosa 1999); (ii) volcano-sedimentary sequences, composed of quartzites, iron formations, pillow lavas and marbles metamorphosed under amphibolite facies, and (iii) the GB intrusives, which occur in the central part of the belt.

The batholith is mainly composed of syenites and monzogranites, with subordinate granites and mafic syenites. Lamprophyric dykes (minettes) also occur (Paim *et al.* 1999). Two main domains can be distinguished (i) multiple intrusions (about 90% of the batholith), and (ii) the so-called late intrusions, which comprise the three massifs Cara Suja (200 km²), Cerafina (120 km²) and Estreito (290 km²). Based on structural data, Rosa *et al.* (1996) propose that the GB was emplaced into a *pull-apart* system.

PETROGRAPHY AND GEOCHEMISTRY The multiple and late intrusions are mainly composed of leucocratic and porphyritic rocks and display similar mineralogy and texture. The magmatic crystallization sequence is: accessory minerals (zircon, apatite, monazite, opaque minerals) ⇒ diopside ⇒ plagioclase ⇒ alkali feldspar

⇒ (amphibole, mica, perthitic feldspar) ⇒ quartz ⇒ titanite. Magmatic crystallization has occurred in two stages: a first stage at low fO_2 , high P_{H_2O} , and transsolvus conditions, accompanied by accessory mineral, feldspar and clinopyroxene crystallization, was followed by a second stage at hypersolvus conditions, characterized by amphibole and feldspar crystallization. The titanite + magnetite + quartz association present at the second stage indicates highly O₂ between the NiNiO and HM buffers.

Geochemical data show that the GB rocks are Si-saturated to -oversaturated, alkalic and metaluminous. The K₂O/Na₂O ratio is always greater than unity, being higher in the mafic members, which can be classified as ultrapotassic (Foley *et al.* 1987). Ba (<8,000 ppm) Sr (<6,000 ppm) and Rb (<940 ppm) are strongly enriched in these rocks. Cr (50 to 700 ppm), Ni (80 to 270 ppm) and Mg# (0.20 to 0.77) are relatively high for common syenitic rocks. Chondrite-normalized REE patterns show strongly fractionated LREE with none or only small negative Eu anomaly. The tectonic discrimination diagrams of the GB rocks as well as their geological context point to a subduction-related and post-collisional setting (Rosa 1999).

U-Pb AND Pb-Pb ZIRCON GEOCHRONOLOGY The U-Pb single zircon analyses were performed at the Swiss Federal Institute of Technology - ETH (Zurich) laboratories (Rosa 1999). The single zircon evaporation Pb-Pb analyses were carried out by Sachet and Macambira (1998) at the Laboratory for Isotope Geology (Para-Iso), Para Federal University, Brazil. The U decay constants are those recommended by IUGS (Steiger and Jäger 1977), and errors are given at the 95% confidence level. The analyzed zircon populations were dominated by brown to light-brown crystals, most of them showing cracks and abundant inclusions.

Single-zircon U-Pb results by isotope dilution techniques The results of six zircon analyses from a sample representative of the multiple intrusions in the Guanambi Region is shown in Fig. 2A. The discordant data points plot on or near a discordia line defining an upper concordia intercept at 2054 ± 8/-6 Ma (MSWD= 7.2, ext. error). Four zircons each were analyzed for three representative members of the late intrusions (Fig. 2A). Three out of four discordant data points from the Cara Suja massif give an upper intercept age of 2053 ± 3 Ma (MSWD= 4.6, int. error), with the fourth data point being displaced to the left of the line. The results for the Cerafina massif show a tight concordant cluster yielding a ²⁰⁷Pb/²⁰⁶Pb age of 2050.4 ± 1.4 Ma (MSWD= 0.67, int. error). The ²⁰⁷Pb/²⁰⁶Pb ages obtained for the four concordant to near concordant data points from the Estreito massif display scatter in excess of their analytical uncertainties. Interpreting the oldest result (2061 ± 5 Ma) as potentially being affected by inheritance, the remaining three grains give a weighted mean ²⁰⁷Pb/²⁰⁶Pb age of 2054 ± 3 Ma (MSWD= 1.64,

1 - Grupo de Petrologia Aplicada - CPGG - Dep. de Geoquímica - IGEO - UFBA, Rua Caetano Moura, 123. CEP:40.210-340, Salvador-Bahia, Brasil (e-mail: lourdes@cpgg.ufba.br)

2 - Swiss Federal Institute of Technology (ETH), Zurich, Switzerland

3 - UMR 6524 CNRS, Université Blaise Pascal, Clermont-Ferrand, France

4 - Laboratorio de Geologia Isotópica (Para-Iso), Universidade Federal do Para-Belem, Brasil




Figure 1 - (A) - Location map of Urandi-Paratinga mobile belt (UPMB) in Bahia. (B) - Simplified geological map of the UPMB. Phanerozoic Cover (1), Meso-Neoproterozoic Covers (2), Guanambi Batholith (3), Cara Suja Massif (3A), Ceraima Massif (3B), Estreito Massif (3C), Volcano-Sedimentary Sequence (4), Santa Izel Complex (5), Bahia (BA), Minas Gerais (MG).

int. error), similar to the age of the other two massifs. The combined results suggest an age of 2.050-2.055 Ga for the crystallization of the GB (2.05-2.06 Ga, if the upper error boundary of the zircons from the Guanambi Region is taken into account). The somewhat disturbed data patterns obtained for three out of the four zircon populations analyzed do not permit to resolve potential time differences for the emplacement of the multiple and late intrusions.

Single-zircon Pb evaporation dating Four crystals from a sample representative of the multiple intrusions (Guanambi Region) analyzed by the stepwise Pb evaporation technique (Kober 1986) yield a mean $^{207}\text{Pb}/^{206}\text{Pb}$ age of 2046 ± 10 Ma (Rosa 1999). Similar ages were obtained for members of the late intrusions, where one zircon from the Ceraima massif shows an age of 2049 ± 2 Ma and four zircons from the Estreito massif have a mean age of 2041 ± 2 Ma (Rosa 1999). Apart from the younger age obtained for the latter, the results are in support of the -2.05 Ga crystallization ages obtained by U-Pb techniques for the GB lithologies.

Rb-Sr AND Sm-Nd ISOTOPIC SIGNATURES Rb, Sr, Sm and Nd whole-rock contents and Sr and Nd isotopic compositions were measured by isotope dilution techniques at the Université Blaise Pascal (Clermont-Ferrand) using a Cameca THN-206 mass spectrometer (Rosa 1999). Uncertainties for $^{147}\text{Sm}/^{144}\text{Nd}$ and $^{87}\text{Rb}/^{86}\text{Sr}$ are 0.5% and 2%, respectively. Initial $^{87}\text{Sr}/^{86}\text{Sr}$, $^{143}\text{Nd}/^{144}\text{Nd}$ and ϵ_{Nd} were computed adopting an age of 2.05 Ga based on the U-Pb results.

Initial $^{87}\text{Sr}/^{86}\text{Sr}$ in the Guanambi Region ranges from 0.705 to 0.707. Similar values of 0.704-0.707 obtained for the late intrusions (Tab. 1) are indicative of a common source. The ϵ_{Nd} values of the GB rocks are strongly negative (-7.4 to -10.6) and imply a LREE-enriched source.

In the $^{87}\text{Sr}/^{86}\text{Sr}$ vs ϵ_{Nd} diagram (Fig. 2B), the GB rocks group tightly in the enriched-source sector. Two contrasting explanations must be considered for this particular pattern: (i) crustal contamination of the GB magma by assimilation of basement rocks, or (ii) derivation from an enriched mantle source. Absence of correlation between initial $^{87}\text{Sr}/^{86}\text{Sr}$ and $1/\text{Sr}$ suggests that crustal contamination was not important. Similarly, there is no correlation between $^{87}\text{Sr}/^{86}\text{Sr}$ and SiO_2 or Al_2O_3 . High Mg#, Ni, and Cr as well as the ubiquitous presence of lamprophyric rocks argue against significant crustal input. We therefore conclude that the chemical and isotopic signature of the GB is inherited from an enriched mantle source.

DISCUSSION AND CONCLUSIONS The 2.05 Ga crystallization age of the GB rocks marks the last magmatic activity in the UPMB region. It further documents a time gap of 0.5 Ga between the emplacement of the syenites of the UPMB and those of the SCMB located in the eastern part of Bahia, which gave a U-Pb zircon age of 2.1 Ga (Conceição 1993). The time interval may reflect changes in the structural regimes of the Paleoproterozoic belts, because, in contrast with the GB rocks, the SCMB syenites are strongly deformed, elongated in a N-S direction and are interpreted as dikes emplaced in a syn-collisional system (Conceição *et al.* 1997), while the GB has intruded a post-collisional environment (Rosa *et al.* 1996).

Several lines of evidence suggest that an enriched mantle source provides the best explanation for the chemical and isotopic signature of the GB rocks. The existence of a Paleoproterozoic enriched mantle in Bahia has already been suggested by Conceição *et al.* (1997) in order to explain the genesis of the Paleoproterozoic syenites of the SCMB. Similarly, Ferreira *et al.* (1997) and Guimarães and da Silva Filho (1998) suggest a metasomatized mantle source for the Neoproterozoic syenitic (NS) Triunfo and Bom Jardim plutons located in the Borborema Province (also in northeastern Brazil). The initial $^{87}\text{Sr}/^{86}\text{Sr}$ ratios of the SCMB and of the Bom Jardim syenites are similar those of the GB rocks (Fig. 2B), whereas ϵ_{Nd} is more positive (0 to -6) in the SCMB syenites and more negative (-10 to -18) in the NS.

Partial melting of an enriched mantle as a source for potassic magmas has been put forward by Wyllie & Sekine (1982). According to these authors, mantle metasomatism leading to growth of phlogopite can occur in the root of subduction zones (~ 100 km) as a result of interaction between cool hydrous siliceous magmas rising from the subducted oceanic crust and the hotter overlying mantle. The nature and composition of the fluids originating from a subduction process are rather complex topics (Wyllie and Sekine 1982, Conticelli and Peccerillo 1992, Ferreira *et al.* 1997). There is, however, some general agreement that the fluids are generated by selective dehydration of the subducting oceanic crust, especially of serpentine and pelagic sediments (clay minerals + carbonates \gg pelites).

Based on Nd model ages, Ferreira *et al.* (1997) and Guimaraes and Da Silva Filho (1998) have argued that mantle metasomatism in the Boborema Province has occurred during the Paleoproterozoic (2.4-2.0




Figure 2 - (A) - U-Pb concordia diagrams for the Guanambi batholith. (B) - Initial $^{87}Sr/^{86}Sr$ vs e_{Nd} for the Guanambi batholith compared to other occurrences of potassic and ultrapotassic rocks. Data are from Conticelli and Peccerillo (1992), Conceicao (1997), Ferreira et al. (1997), Guimaraes and Da Silva Filho (1998).

Table 1 - Rb-Sr and Sm-Nd isotopic data for the Guanambi batholith. Multiple intrusions (MI), Cara Suja (CS), Ceraíma (CE), Estreito (ES).

Sample	Locality	Rb (ppm)	Sr (ppm)	$^{87}Rb/^{86}Sr$	$^{87}Sr/^{86}Sr$	$(^{87}Sr/^{86}Sr)_i$	Sm (ppm)	Nd (ppm)	$^{147}Sm/^{144}Nd$	$^{143}Nd/^{144}Nd$
996	MI	127	1009	0.3642	0.716735	0.7060	24.60	186.2	0.0799	0.510591
1002	MI	135	883	0.4447	0.719331	0.7062	27.03	177.4	0.0922	0.510725
1008	MI	229	656	1.0112	0.736326	0.7065	24.20	168.6	0.0868	0.510686
1128	MI	146	985	0.4287	0.719936	0.7073	26.07	172.8	0.0913	0.510778
1139	MI	158	726	0.6307	0.724695	0.7061	25.99	168.2	0.0935	0.510781
1141	MI	236	1259	0.5414	0.723446	0.7075	21.95	141.4	0.0939	0.510831
1216	MI	191	794	0.6970	0.725691	0.7051	19.58	133.8	0.0885	0.510698
1146	MI	193	385	1.4511	0.746839	0.7040	21.58	188.0	0.0694	0.510542
1160	MI	271	1236	0.6337	0.724054	0.7053	17.83	107.3	0.1005	0.510906
974	CS	340	1159	0.8553	0.72939	0.7060	25.31	161.9	0.0946	0.510877
982	CS	330	454	2.1007	0.76861	0.7041	8.04	54.1	0.0898	0.510737
983	CS	261	560	1.3463	0.74581	0.7066	24.79	186.1	0.0806	0.510591
985	CE	188	1519	0.3576	0.715466	0.7049	36.30	216.3	0.1015	0.510819
1170	CE	175	6192	0.0818	0.708121	0.7057	25.12	148.1	0.1026	0.510847
1200	CE	197	1762	0.3226	0.71525	0.7057	29.80	177.0	0.1016	0.510846
1009	ES	179	1347	0.3837	0.716717	0.7054	6.94	46.6	0.0902	0.510659
1014	ES	237	1493	0.4590	0.717659	0.7041	20.17	127.5	0.0957	0.510762
1228	ES	157	477	0.9520	0.735404	0.7073	28.53	288.4	0.0598	0.510281

Ga). Similar chemical and isotopic characteristics suggest that this interpretation also applies to the GB rocks, which we view as products of fractional crystallization of a lamprophyric magma generated by melting of a Paleoproterozoic enriched mantle, as indicated by Conceição *et al.* (1997).

The bulk of evidence available so far suggests that a lithospheric mantle region enriched during a Paleoproterozoic event has repeatedly served, over an extensive time interval of at least 1.5 Ga, as a source for alkaline magmatism in northeastern Brazil and in Bahia State. Independent of the tectonic regime, this mantle source has been

particularly productive during the Paleoproterozoic in generating the vast volumes of syenitic rocks of the mobile belts.

Acknowledgments This paper was supported by CAPES (Process: BEX 3065/95-13 and BEX 2521/97-4), Companhia Baiana de Pesquisa Mineral (CBPM) and CNPq (Process: 521592/97-6). We thank C.G. Perreira Santos for drafting the map and two anonymous referees of RBG for their suggestions, which improved the manuscript. This is the contribution number 72 of GPA-CPGG-UFBA and is a contribution to IGCP-426.

References

- Conceição H. 1993. Petrology of the syenites from Salvador-Curaçá Mobile Belt (Bahia-Brazil): geodynamic significance. *An. Acad. Bras. Ci.*, 65(1):17-32.
- Conceicao H., Martin H., Rosa M.L.S., Conceição R.V., Sabate P. 1997. Alkali-potassic magmas generated by partial melting of an enriched mantle source: example of the lower Proterozoic plutonism, State of Bahia, NE Brazil. II ISGAM, *Ext. Abstr.*, 1 OS-109.
- Conticelli S. and Peccerillo A. 1992. Petrology and geochemistry of potassic and ultrapotassic volcanism in central Italy: petrogenesis and inference on the evolution of the mantle sources. *Lithos*, 28(3-6):221-240.
- Ferreira V.P., Sial A.N., Long L.E., Pin C. 1997. Isotopic signature of Neoproterozoic to Cambrian ultrapotassic syenitic magmas, Northeastern Brazil: evidence for an enriched mantle source. *International Geology Review*, 39:660-669.
- Foley S.F., Venturelli G., Green D.H., Toscani L. 1987. The ultrapotassic rocks: characteristics, classification and constraints for petrogenetic models. *Earth Sci. Rev.*, 24:81-134.
- Guimaraes I.P. and Da Silva Filho A.F. 1998. Nd- and Sr-isotopic and U-Pb geochronologic constraints for evolution of the shoshonitic Brasileiro Bom Jardim and Toritama Complexes: evidences for a Transamazonian enriched mantle under Borborema Tectonic Province, Brasil. *International Geology Review*, 40:500-527.
- Kober B. 1986. Whole-grain evaporation for $^{207}\text{Pb}/^{206}\text{Pb}$ -age-investigations on single zircons using a double-filament thermal ion source. *Contrib. Mineral. Petrol.*, 93:482-490.
- Paim M.M., Rosa M.L.S., Conceição H., Barreto Santos E., Bastos Leal L.R. 1999. Aspectos geoquímicos dos lamprofiros (minettes) associados ao maciço sienítico de Cara Suja - Sudoeste da Bahia. V Congresso de Geoquímica dos Países de Língua Portuguesa & VII Congresso Brasileiro de Geoquímica, Porto Seguro-BA, Anais:566-568.
- Rosa M.L.S. 1999. *Geologia, Geocronologia, Mineralogia, Litogeoquímica e Petrologia do Batólito Monzo-Sienítico Guanambi-Urandi (SW-Bahia)*. Instituto de Geociências, Universidade Federal da Bahia, Salvador, Tese de Doutorado, 186p.
- Rosa M.L.S., Conceicao H., Paim M.M., Barreto Santos E., Alves Da Silva EC., Leahy G.A.S., Bastos Leal L.R. 1996. Magmatismo potássico-ultrapotássico pos a tardi-orogênico (associado a subducção) no oeste da Bahia: Batólito Monzo-Sienítico de Guanambi-Urandi e os Sienitos de Correntina. *Geochim. Brasil*, 10(1):027-042.
- Sachet C.R. and Macambira M.J.B. 1998. *Estudo geocronológico pelo método de evaporação de Pb em zircões do Batólito Guanambi-Urandi - BA*. Relatório Técnico, 8p. (Inédito).
- Steiger R.H. and Jager E. 1977. Subcommittee on geochronology. Convention on the use of decay constants in geo- and cosmochemistry. *Earth Planet. Sci. Letters*, 36:359-362.
- Wyllie P.J. and Sekine T. 1982. The formation of mantle phlogopite in subduction zone hybridization. *Contrib. Mineral. Petrol.*, 79:375-380.

Contribution IGC-096
Received March 1, 2000
Accepted for publication April 6, 2000