

U–Pb and Sm–Nd geochronology of amphibolites from the Curaçá Belt, São Francisco Craton, Brazil: Tectonic implications

Luiz José Homem D’el-Rey Silva ^{a,*}, Elton Luiz Dantas ^b, João Batista Guimarães Teixeira ^c,
Jorge Henrique Laux ^b, Maria da Glória da Silva ^{c,d}

^a Universidade de Brasília — Instituto de Geociências

^b Laboratório de Geocronologia, Universidade de Brasília, Campus Darcy Ribeiro, Asa Norte, 70.910-900, Brasília — DF, Brazil

^c Grupo de Metalogênese, Centro de Pesquisa em Geofísica e Geologia, Universidade Federal da Bahia, Campus Universitário de Ondina, Sala 201-C, Salvador, 40170-290, Bahia, Brazil

^d Serviço Geológico do Brasil (CPRM) - Av. Pasteur 404, Urca, 22290-240 Rio de Janeiro — RJ, Brazil

Received 24 May 2006; received in revised form 29 July 2006; accepted 23 November 2006

Available online 16 January 2007

Abstract

The Curaçá terrane is part of the Itabuna–Salvador–Curaçá (I–S–C) Paleoproterozoic orogen in the São Francisco craton, northeastern Brazil, and comprises supracrustal rocks, gneisses of their probable basement, amphibolites, and mafic-ultramafic Cu-bearing bodies (including the Caraíba Cu-Mine), all affected by D₁–D₃ deformation events associated to M₁–M₃ metamorphism under high-T granulite and amphibolite facies, and assisted by G₁–G₃ tonalitic-granodioritic-granitic intrusions. U–Pb and Sm–Nd Thermal Ionization Mass Spectrometry (TIMS) isotopic data from amphibolite, tonalite, and granite, sampled in a well-known outcrop, indicate partial reset and heterogeneous modification of the original isotopic systems, attributable to deformation and metamorphism. The ages obtained from these systems agree with each other, and also with other previously published U–Pb data, and imply that 2.6 Ga is the crystallization age of the protolith of the amphibolite. Together with key structural relationships, they also indicate a 2.08–2.05 Ga interval for M₃ metamorphism, and make even a less precise age (2.2–2.3 Ga) acceptable, as it suggests contamination in the amphibolite with material in a syn-D₂ tonalite crystallized 2248±36 Ma ago. The new data demonstrate the existence of Neoproterozoic fragments of both oceanic and continental crusts and constrain the Archean–Paleoproterozoic development of the Curaçá belt, the I–S–C orogen, and the São Francisco craton.

© 2006 International Association for Gondwana Research. Published by Elsevier B.V. All rights reserved.

Keywords: U–Pb and Sm–Nd isotopic data; Curaçá terrane; São Francisco craton; Archean to Paleoproterozoic tectonics; Caraíba Mine

1. Introduction

Proterozoic mobile belts or terranes of high metamorphic grade are among the most characteristic features of continental shield areas (Windley, 1986; Condie, 1989). These terranes consist of polydeformed/metamorphosed rocks displaying a generally older regional metamorphic banding that may hide the evidence for polycyclic tectonic events recognisable only upon detailed mapping combined with litho-geochemistry (Passchier et al., 1990). Consequently accurate dating of protoliths crystallization and/or deformation and metamorphism in such

belts relies on detailed structural studies of key outcrops combined with geochronological methods (Gromet, 1991).

The Curaçá River Valley terrane is the N–S trending northern segment of the Itabuna–Salvador–Curaçá (I–S–C) Paleoproterozoic orogenic belt that stretches between the Mesoarchean Gavião, Serrinha, and Jequié cratonic blocks, in the central-eastern part of the São Francisco craton, Bahia, eastern Brazil (Fig. 1; Sabaté et al., 1990; Barbosa et al., 1996). The existence of reworked Archean rocks in the Curaçá terrane (Inda and Barbosa, 1978; Lindenmayer, 1981), and across the entire I–S–C orogen, has driven Leite (2002) and Barbosa and Sabaté (2002, 2004) to postulate the participation of a fourth Archean block inside the Paleoproterozoic collision zone. Geodynamic models for the Archean–Paleoproterozoic evolution of the São Francisco craton have been supported by recent

* Corresponding author. Tel.: +55 61 30324907; fax: +55 61 33474062.

E-mail addresses: ldel-rey@unb.br, luizhomemsilva@yahoo.com.br (L.J.H. D’el-Rey Silva).




Fig. 1. a — Summary display of the Gavião, Serrinha and Jequié blocks, and the Itabuna–Salvador–Curaçá (I–S–C) orogen, all within the geographical limits of the Bahia State, and part of the São Francisco craton, Brazil. Numbers in brackets represent T_{DM} model age data for each block (adapted from Barbosa and Sabaté, 2004). b — Simplified geological map of the part of northern Bahia to show the northern part of the I–S–C orogen (= Curaçá terrane), between the Gavião and the Serrinha Archean blocks (combining Figs. 1 in D'el-Rey Silva, 1984, 1985; and in Oliveira et al., 2004). The Caraíba orebody and the area of Fig. 2 are highlighted. Details in text.

isotopic studies (Oliveira et al., 2004; Barbosa and Sabaté, 2004), indicating that the I–S–C orogen and the three Archean blocks preserve distinct Sm–Nd characteristics (Fig. 1a).

Early regional studies (Delgado and Souza, 1975) made the Curaçá terrane well-known because of its high potential for Cu-sulphide mineralization associated with mafic–ultramafic (M–UM) bodies mapped within high metamorphic grade supracrustal rocks (metasediments and metavolcanics), gneisses, migmatites, and metagranitoids. In fact, following the accidental discovery of green copper oxide (malachite) nearly 140 years ago in the site destined to become the Caraíba Mine (Fig. 1b), six other smaller Cu-deposits have been found in the two last decades, among over two hundred small bodies (D'el-Rey Silva, 1984, 1985). Structures record progressive (D_1 – D_3) deformation (Fig. 2; D'el-Rey Silva, 1993) and conditions for the main metamorphism have been estimated around 850°–720 °C (Jardim de Sá et al., 1982; Ackermann et al., 1987), or up to 1000 °C, with pressures varying from 8–10 kbar (Leite, 2002; Leite et al., 2005), all on the basis of several geothermometers and sapphirine-bearing assemblages.

The first U–Pb isotopic data for rocks of the Curaçá belt, all from the surroundings of the Caraíba orebody (Fig. 2), indicated

crystallization ages of: 2580 ± 10 Ma for a sample of norite collected at the Caraíba Mine (U–Pb Sensitive High Resolution Ion Microprobe – SHRIMP – data in Oliveira et al., 2004); 2248 ± 36 Ma for syn- D_2 tonalite in the Caraíba Airport outcrop; and 2051 ± 16 Ma for syn- D_3 granite intrusive in the Caraíba orebody (both U–Pb data, but respectively from zircons and monazite; D'el-Rey Silva et al., 1996), whereas the age of the supracrustals was not directly determined.

This paper presents the results of U–Pb and Sm–Nd integrated studies carried out on samples of amphibolite, tonalite and granite of the Caraíba Airport outcrop, situated ~5 km to the north of the Caraíba orebody (Fig. 2). This outcrop is one of the most visited places in the Curaçá belt exactly because it displays clear evidence for a D_1 – D_3 progressive deformation under M_1 – M_3 metamorphism, providing a well-constrained structural–metamorphic framework for targeting geochronological studies.

2. Regional geology and tectonic setting

The basement of the São Francisco Craton is mostly represented by the Gavião and Serrinha blocks, shown




Fig. 2. Simplified geological map of the area surrounding the Caraíba orebody, highlighting the Curaçá antiform, the open pit, and the land track of the airport. Based on a 1:10,000 scale geological mapping carried out by D'el-Rey Silva (1984, 1985) along 130 km of E–W tracks distributed in 18 lines 500 m apart one from each other, some marked every 100 m, others marked every 50 m, a real frame (omitted for simplicity) opened in the field for geophysical survey. Mapping of the orebody itself was at 1:1,000 scale (see Fig. 3a) whereas the part to the N of the open pit (belonging to DOCEGEO Company) was mapped at 1:5,000 scale (Lindenmayer et al., 1984). See details in text.

respectively to the west and to the east of the Curaçá terrane, in their simplest map expression (Fig. 1b). The Gavião block is mainly composed of gneiss-amphibolite associations, amphibolite-facies tonalite-granodiorite orthogneisses dated at 2800–2900 Ma as well as greenstone belts. The block also encloses a 3200–3400 Ma trondhjemite-tonalite-granodiorite suite (Bar-

bosa and Sabaté, 2004). Its westernmost part is covered by (meta)sediments deformed in the Neoproterozoic Brasiliano orogeny and by Quaternary limestones, whereas the eastern part includes (to the west of the town of Bonfim, Fig. 1b) the Campo Formoso Cr-bearing mafic–ultramafic layered body, a major intrusion that is unconformably overlain by the Jacobina Group,

both of which are intruded by the Late Paleoproterozoic Campo Formoso granite (Silva, 1996; Leite, 2002). According to a summary in Silva (1996) the Jacobina Group is part of a much longer Paleoproterozoic rift-like volcanic-sedimentary basin extending to the boundary between the Gavião and Jequié blocks.

The Serrinha block consists of amphibolite-facies 2900–3500 Ma ortho and paragneisses of granodiorite composition, as well as migmatites and amphibolites (Barbosa and Sabaté, 2004). The western margin is in contact with the Itiúba syenite, whereas the eastern margin encloses Paleoproterozoic greenstone sequences, and arc-like orthogneiss and granites that are indicators of a back-arc basin system within the Serrinha block (Silva, 1996). Farther to the east, this block underlies Neoproterozoic (meta)sediments and, outside of Fig. 1b, Mesozoic sediments of the Tucano basin.

The I–S–C orogen (Fig. 1) is mainly composed of granulite-facies tonalitic and charnockitic rocks with basic-ultrabasic enclaves, as well as supracrustal rocks, in both of its northern and southern segments (Barbosa and Sabaté, 2004). The 2084 Ma old Itiúba syenite separates the 2580 Ma old Curaçá Valley Cu-district to the west, from the 2085 Ma old Jacurici Valley Cr-district to the east (Fig. 1b; all U–Pb SHRIMP age of zircons; Oliveira et al., 2004).

3. Regional geology of the Curaçá high-grade terrane

3.1. Lithostratigraphy

The lithotypes of the Curaçá terrane are divided into the Tanque Novo and Caraíba complexes, both including the Cu-bearing mafic-ultramafic bodies (Fig. 1b). The Tanque Novo Complex comprises Al-rich paragneisses, banded gneisses, calcsilicate rocks, quartzites, iron formations and graphite gneisses, whereas the Caraíba complex comprises mostly enderbite and charnockitic orthogneisses, as well as migmatites.

Most of the regional geology and structural evolution of the Curaçá belt (Lindenmayer, 1981; Figueiredo, 1981; Mandetta, 1982; Gáal, 1982; Hasuy et al., 1982; Jardim de Sá et al., 1982; D'el-Rey Silva, 1984, 1985), published at the same time or soon after the start-up of the Caraíba Mine operation, have shown the Curaçá terrane as formed of three main lithostratigraphic units also recorded in the vicinities of the Caraíba orebody (Fig. 2), the area submitted to the most detailed mapping carried out in the Curaçá belt.

The Tanque Novo Complex is a supracrustal sequence consisting mostly of quartzfeldspathic gneiss with thin intercalations of amphibolites, cordierite-sillimanite-garnet paragneiss, oxide-facies banded iron formation, calcsilicate rocks, anhydrate-bearing marbles, forsterite-marbles and quartzites, as well as gneisses supposed to belong, at least in part, to the crystalline basement. The second unit consists of Cu-poor or simply barren gabbros, gabbronorites, leucogabbros, and hyperites, together with Cu-rich hypersthenites, melanorites, and norites. A vast amount of geology and structural data collected regionally, and also on surface and in underground sites of the Caraíba Mine and other small bodies, drove

Lindenmayer (1981), Lindenmayer et al. (1984), D'el-Rey Silva (1984, 1985), and lately Mayer and Barnes (1996) and D'el-Rey Silva et al. (1988, 1994, 1996) to state that the Cu-bearing rocks are sill-like structures. It is noteworthy that the Caraíba orebody is a Cu-only, chalcopyrite and bornite-rich deposit and associated with deep ocean rocks such as banded iron formation and evaporites (anhydrate) forsterite-marble. The third unit (the Caraíba complex) mostly comprises migmatitic gneisses and syntectonic intrusions (G_1 and G_2) generally consisting of grey tonalites and granodiorites. Reddish-pink coloured, K-rich (G_3) granites intruded all lithotypes and developed in association with D_3 (Fig. 2).

3.2. Structures and metamorphism

The Curaçá belt has been affected by a D_1 – D_3 progressive deformation assisted by large volumes of syntectonic G_1 – G_3 granitoid intrusions. Associated metamorphisms M_2 (granulite facies) and M_3 (amphibolite facies) are quite evident everywhere, but evidence for a M_1 amphibolite facies metamorphism is solely preserved in paragneisses and amphibolites that occur as dm- to m-scale boudins or xenoliths entrained in G_2 intrusions across the belt (Jardim de Sá et al., 1982; D'el-Rey Silva, 1984, 1985).

Jardim de Sá et al. (1982) first reported M_1 amphibolite-facies metamorphism on the basis of: 1 — Migmatites affected by D_2 and D_3 events; 2 — Biotite inclusions (M_1) found in crystals of orthopyroxene (M_2) in paragneisses; and 3 — An opx-rich border surrounding layers and boudins of mafic rocks (amphibolites) in paragneisses, all affected by F_3 folds associated with M_3 metamorphism (their Fig. 4H). Whereas these observations are respectively from outcrops to the south and north of Caraíba (the two latter in the Tanque Novo Sequence, Surubim area; Fig. 1b), the existence of M_1 metamorphism in the Caraíba area (Fig. 2) is demonstrated by several inclusions of circular crystals of green hornblende in hypersthene from the opx-rich border of mafic boudins in the Airport outcrop (photograph 45, page 153 of D'el-Rey Silva's MSc thesis, 1984). Hypersthene-free amphibolites have been largely documented across the Curaçá Valley (modal compositions in page 99 of Lindenmayer, 1981). M_2 mineral assemblages include cordierite-sillimanite-garnet-biotite in paragneisses, and andesine plus hypersthene (strongly pleochroic and rich in Al_2O_3) in commonly granoblastic gabbroic rocks, even the Cu-bearing bodies (Lindenmayer, 1981). M_3 metamorphism resulted in the transformation of hypersthene into hornblende and biotite, in F_3 hinges (photographs 50 and 51, D'el-Rey Silva's MSc thesis) and is characterized by biotite, hornblende, quartz and feldspar assemblages (Lindenmayer, 1981).

Due to D_1 – D_3 evolution, the older metamorphic banding (S_1) and intrafolial folds F_1 are affected by two folding phases (F_2 and F_3), so that S_1 is mostly a composite structure where evidences for M_1 is commonly absent, but M_2 and M_3 assemblages dominate. S_1 appears in most outcrops as a continuous banding traceable for 10 m, or more, and consisting of cm- to dm-thick bands of dark, more mafic (hornblende,

biotite, pyroxene) minerals, and bands of light, more felsic minerals (feldspars and quartz, mostly). S_1 may also appear as mm- to cm-thick, generally continuous in ≤ 1 m-scale, or somehow diffuse banding in metabasic rocks and amphibolites such as the ones in the studied outcrop. S_1 is also a banding parallel to S_0 defined by the intercalation of 10 cm- to 50 cm-thick layers of acid gneiss (possibly meta-arkose and/or meta-rhyolite), amphibolite, pelitic paragneiss, marble, banded iron formation, and calcsilicate rocks.

F_1 folds are generally of 10 cm-scale, rootless and intrafolial relative to S_1 (S_0), whereas the F_2 folds (and foliation S_2) are mostly seen in outcrops of gneisses and orthogneisses, together with examples of 10–100 cm-sized patterns indicative of Ramsay's (1967) Type 2 of $F_2 \times F_3$ fold interference pattern (Jardim de Sá et al., 1982; D'el-Rey Silva, 1984) but could be better understood only after detailed mapping of the Caraíba orebody and surrounding area (next section). F_3 and S_3 are also very common features in the Curaçá terrane. F_3 folds are non-cylindrical, cm- to km-sized, generally tight, asymmetric, E-verging, and their axial planes trend nearly N–S and dip 70° – 75° W. The F_3 axes (B_3) plunge generally less than 30° , either in a northerly or southerly direction. S_3 is a typical mineral foliation mostly defined by biotite, hornblende, and strongly flattened quartz/feldspars, but the non-micaceous minerals display a prolate geometry and define a penetrative L_3 mineral stretching lineation parallel to B_3 . Granites G_3 are early- to late- D_3 (Fig. 2) and intruded sub-parallel to the axial trace of F_3 folds. They most commonly display the S_3 foliation, and represent typical lithotypes due to M_3 metamorphism (Fig. 2). The Curaçá antiform (Delgado and Souza, 1975; Gáal, 1982) is a D_3 structure, the hinge of which plunges to the N or to the S (reference to the Surubim area) and is the site for most of the mafic–ultramafic bodies (Fig. 1b). Also in the Surubim area, Jardim de Sá et al. (1982) and Hasuy et al. (1982) described 1 m- to 10 m-sized structures of Type 1 (domes and basins) interference pattern between the regional F_3 folds and gentle-open, very local F_4 folds that have no influence in the geometry and spatial distribution of the layers.

4. The Airport outcrop: key geological features preserved in an extremely deformed area

The Airport outcrop is a flat-lying exposure of tonalitic orthogneiss with an amœboid surface area of nearly 1000 m², and lies in the northern end of the land track of the Mine airport, and is situated in the western limb and close to the hinge of the Curaçá antiform, in a similarly structural position as the Caraíba orebody (Fig. 2). The orthogneiss also encloses tens of boudins of amphibolite, the whole set intruded by sub-vertical, sheet-like bodies of pink granites, and some quartz-feldspar veins. The outcrop itself displays important clues for understanding the critical role of early E–W trending structures, such as the F_2 folds, in the evolution of the Curaçá terrane (next section), but a complete understanding of the importance of the outcrop, and the importance of dating the amphibolites (see Discussion) all demand an overview on the geology of the Caraíba orebody, where such F_2 folds were fully

mapped (Fig. 3a), and on the tectonic conditions prevailing in the I–S–C orogen.

D_3 flattening was so strong and melt-assisted that the real geometry of some of the 10 cm- to 1 m-scale $F_2 \times F_3$ fold interference patterns seen in the Curaçá terrane is often obscured by disrupted hinges, melts injection, and/or extreme ductile thinning. However, the large amount of data from detailed mapping of the orebody and its surroundings (1978–1984; D'el-Rey Silva, 1984, 1985) and from mine operation in the following 18 years (D'el-Rey Silva et al., 1988, 1994, 1996; D'el-Rey Silva and Oliveira, 1999) allows a picture to be constructed of the Caraíba Cu-deposit within the core of a 1000 m-high, tight, non-cylindrical F_3 synform gently-moderately plunging to the S (a parasite fold of the Curaçá antiform). It acquired the shape of a N–S trending mushroom (Figs. 2 and 3a), due to the interference of the F_3 synform with several F_2 folds, so that the E–W trending and originally sub-horizontal F_2 axis (B_2) appears sub-vertical in the upper half of the orebody (Fig. 3b), and the pre- D_3 attitude of the layers also forces F_3 fold axes to plunge steeply to the N solely in the central and very inner part of the mushroom. Thus, the orebody consists, to the S of section 32 and to the N of section 37 (Fig. 3b), of a mineralized layer affected by simple synform, whereas between sections 32 to 37 (the central and richer part) it displays repetitions of the same layer due to several F_2 hinges affected by the F_3 synform (Fig. 3c and inset).

The importance of the Airport outcrop and the Caraíba orebody rests in the preservation of key evidence exactly where the I–S–C orogen was squeezed to its narrowest map expression (Fig. 1a–b). Similar D_1 – D_3 structures and their field relationships are found further south, where the orogen is wider (in the Ipirá-Lajedinho area, Fig. 1a). In that area, largely protected by the northern margin of the Jequié block, abundant, m- to dm-scale, E–W trending F_2 folds associated with mapable, ductile, D_1 -related lateral ramps, N–S trending D_3 transcurrent faults, as well as $F_2 \times F_3$ interference patterns, some as large as the Caraíba mushroom, justifies interpreting the orogen as the result of a long-lived, oblique collision-related sinistral transpressional orogen (D'el-Rey Silva, 1993; Barbosa, 1996), and the intensity of shortening peaks around Caraíba for two reasons: 1 — the area surrounding Caraíba lies 300 km away from the Jequié block; and 2 — intrusion of the D_2 -related syenite most likely added to make even stronger the E–W compression responsible for the transposition of any pre- D_3 structures that existed in the Caraíba region.

The tonalite in the Airport outcrop is yellowish-grey, well-banded, and composed of oligoclase-andesine, hornblende, biotite, orthopyroxene, clinopyroxene, minor microcline and quartz, with garnet, magnetite, apatite, and zircon as accessories. It also encloses cm- to dm-thick and >2 m-long bands of acid composition, as well as <1 m-sized lenses of non-porphyrific granitoid, all aligned parallel to S_2 , this one a quite homogeneous granulite-facies metamorphic banding consisting of ≤ 1 cm-thick bands of flattened crystals of feldspar, quartz, minor biotite, intercalated with ≤ 1 cm-thick bands of pyroxene, hornblende, plus biotite, and affected by numerous, 0.5–2.0 m-scale, very tight F_3 folds that plunge $\leq 25^\circ$ to the S and are




Fig. 3. a — Simplified geological map of the Caraíba orebody with indication of the S₁ foliation, minor D₃ structures, as well as the axial trace of the main F₂ and F₃ folds. The walls of each of the four upper benches of the open pit were mapped (1978–1984) in a 1:1,000 scale, simultaneously with the opening of bench 395 m, the lower one at that time, so the orebody could be tracked in great detail as new exposures were available every 10–20 m along strike, blast after blast. The geometry of the orebody match entirely the Cu-grade iso-contour map obtained on each bench by the plot of thousand data on the Cu-grade from drill holes of a 5 m-square grid used for blasting. The eastern limb of the mushroom structure occurs only underground, from cross section 35 to the N, and has been projected on surface as a modification here introduced in the original map by D'el-Rey Silva (1984, 1985); b — A 3-D schematic representation of the Caraíba mushroom (from D'el-Rey Silva et al., 1988); c — The overall pattern of F₂ × F₃ interference for cross section 36 (see also the inset) is typical for sections 32–37 (the central part of the orebody). Diamond drill holes are indicated for different years. Adapted from D'el-Rey Silva et al. (1996) and D'el-Rey Silva and Oliveira (1999). See text for details.

associated with a sub-vertical, N–S striking, and penetrative mineral foliation (S_3) marked mostly by biotite, but also by hornblende plus quartz and feldspar. S_3 can be seen cross-cutting at very low angle the S_1 banding in the boudins.

Only after the detailed mapping (Fig. 4), can the boudins be seen to define nearly E–W trending trails affected by F_3 folds, revealing a more complete story: during D_2 the G_2 tonalite intruded continuous layers of E–W trending S_1 -foliated




Fig. 4. Simplified geological map of the Caraiba Airport outcrop to show the site of collection of the samples used in this study. B_3 fold axis plunges $\leq 25^\circ$ to the S, whereas planar structures such as S_1 in the amphibolite boudins, S_2 , and S_3 , all dip sub-vertically, unless in the very hinge of the F_3 folds. G_3 granite intrusions cut across some of the F_3 hinges. Based on the original lithostructural mapping at 1:100 scale carried with the aid of a plane table and alidade by L.J.H. D'el-Rey Silva and J.A.C. de Morais (D'el-Rey Silva, 1984, 1985). Figures indicative of $F_2 \times F_3$ interference pattern (some observed, other highly suggested) are indicated in the areas surrounded by ellipses. Details in text.

amphibolite. The amphibolite layers underwent boudinage while the tonalite deformed by ductile flow and acquired the granulite-facies S_2 banding. During D_3 the boudins rotated nearly 90° around a vertical axis, so their longer horizontal axis, originally oriented nearly E–W, became systematically oriented N–S, the boudin ellipses entrapped in the F_3 hinges (Fig. 5a, b, e)

and also underwent D_3 shortening. The boudins systematically nested in the F_3 hinges, with S_1 and the long axis both parallel to S_3 , plus the abundant $F_2 \times F_3$ folding interference patterns in the tonalite, or highly suggested in the amphibolites (Fig. 4), altogether strongly support interpreting that: 1 — S_2 formed sub-parallel to S_1 ; 2 — the G_2 tonalite is a syntectonic intrusion; 3 —




Fig. 5. Features of the Airport outcrop, as seen in four view to the S photos (a, b, c, e) and in two photos from thin-sections (d, f). Several boudins (a) define a E–W trending layer affected by F_3 folds, in the central part of the outcrop. The longer axis of the boudins parallels the hammer's handle that points to the S. Technician R. Belo is partially shown on the upper left corner, beside the plane table and alidade used for detailed mapping; b — A 10 cm-thick acid band and the grey tonalite affected by an F_3 fold around the northern margin of the largest boudin (dotted line) in the outcrop. The traces of S_1 and AP_3 are parallel one to each other; c — Relationships between the F_3 folded tonalite and the eastern side of the boudin source of sample 1A. Note the <50 cm-wide felsic pegmatoid with over 1 cm-big crystals of orthopyroxene along the amphibolite-tonalite contact; d — Detail of a thin section from sample collected at the boudin-tonalite contact (in c). Large crystals of hornblende (Hb) dominate in the left half of the picture, and occur together with plagioclase and some crystals of orthopyroxene (Opx), whereas the latter dominates in the right half, together with larger crystals of plagioclase. Opx crystals grew at the expenses of the Hb crystals along the margin of the boudin, as also demonstrated by circular crystals of Hb included in Opx (D'el-Rey Silva, 1984, 1985); e — Detail of a F_3 fold affecting S_2 around the head of a boudin. Note the traces of S_1 (inside the boudin) and S_3 both parallel to the boudin's long axis; and f — Picture of part of a thin-section beside a F_3 hinge. Note crystals of pyroxene (some marked by *) and plagioclase along S_2 , as well as ribbons of biotite and flattened crystals of plagioclase defining the S_3 foliation.

Table 1
Summary of the U–Pb data from thirteen fractions of zircons from 10 kg of the amphibolite

U (ppm)	Pb (ppm)	Pb206 Pb204 (obs)	Pb207* U235	(pct)	Pb206* U238	(pct)	Correl. coeff. (rho)	Pb207* Pb206*	(pct)	Pb206* U238 Age (My)	Pb207* U235 Age (My)	Pb207* Pb206* Age (My)	(My)
145	76	1596	7.56206	0.57	0.37703	0.571	0.99019	0.145466	0.08	2062	2180	2293	1.4
196	71	862	6.36684	0.83	0.360684	0.791	0.94863	0.128025	0.263	1985	2027	2071	4.6
49	17	448	5.38241	1.24	0.311302	1.200	0.96828	0.125399	0.309	1747	1882	2034	5.5
205	32	297	1.40039	1.84	0.142056	1.650	0.90839	0.071497	0.769	856	889	971	16
201	74	815	6.32118	0.66	0.360535	0.636	0.966734	0.127160	0.168	1984	2021	2058	3
174	65	857	6.42752	0.68	0.363910	0.663	0.975212	0.128100	0.150	2000	2036	2072	2.6
546	202	604	5.99644	0.45	0.344294	0.431	0.959408	0.126317	0.128	1907	1075	2047	2.3
330	108	817	7.42198	1.04	0.287785	1.040	0.997556	0.1870470	0.729	1630	2163	2716	1.2
90	13	916	2.0111	1.02	0.145027	1.020	0.925758	0.1005740	0.383	873	1119	1634	7.2
577	116	7745	3.55528	0.5	0.201059	0.496	0.991762	0.1282480	0.063	1181	1539	2074	1.1
313	132	1116	7.57359	0.35	0.362173	0.349	0.994050	0.1516650	0.004	1992	2181	2364	0.65
206	100	582	8.57972	0.7	0.392694	0.633	0.906718	0.1584590	0.294	2135	2294	2439	5
309	84	383	2.9353	1.02	0.199033	0.994	0.977454	0.1069610	0.215	1170	1391	1748	4

D₁–D₃ deformation was progressive; and 4 — lithological layers, as well as S₁ and S₂ foliation, may all have had an E–W trend somewhere in the Curaçá terrane, before D₃ deformation. Thus, it is possible that the amphibolite layers were affected by E–W trending, 10–100 m sized, isoclinal F₂ folding, in the hinges of which the G₂ tonalite intruded. The double row of boudins in the central part of the outcrop (Fig. 4) suggests a 10 m-sized F₂ × F₃ boomerang-like Type 2 interference pattern. The strength of interpreting and the validity of the U–Pb diffusion technique age data (next section) rely in the validity of these structural and metamorphism relationships demonstrated above, and also in the fact that such ages agree with others obtained for the same rocks, using more precise techniques, and are subject of a detailed discussion at the end of this paper.

5. Geochronology

5.1. Sample population and analytical procedures

The isotopic data derive from eleven samples of rocks collected in the Airport outcrop (Fig. 4), then treated and analyzed in the Laboratory of Geochronology — University of Brasília. Zircon concentrates were extracted using conventional gravimetric (DENSITEST®) and magnetic (Frantz isodynamic separator) techniques applied to ca. 10 kg rock samples collected in the largest boudin in the outcrop. Final purification was achieved by handpicking using a binocular microscope. For the conventional U–Pb analyses, fractions were dissolved in concentrated HF and HNO₃ (HF:HNO₃=4:1) using micro-capsules in Parr-type bombs. A mixed ²⁰⁵Pb–²³⁵U spike was used. Chemical extraction followed standard anion exchange technique, using Teflon micro columns, following procedures modified from Krogh (1973). Pb and U were loaded together on single Re filaments with H₃PO₄ and Si gel, and isotopic analyses were carried out on a Finnigan MAT-262 multi-collector mass spectrometer equipped with secondary electron multiplier — ion counting. Procedure blanks for Pb, at the time of the analyses, were better than 20 pg. PBDAT (Ludwig, 1993) and ISOPLOT-Ex (Ludwig, 2001a) were used for data reduction and age calculation. Errors for isotopic ratios are 2σ.

The Sm–Nd isotopic analyses followed the method described by Gioia and Pimentel (2000): whole rock powders (ca. 50 mg) were mixed with ¹⁴⁹Sm–¹⁵⁰Nd spike solution and dissolved in Savillex capsules. Sm and Nd extraction of whole-rock samples followed conventional cation exchange techniques, using teflon columns containing LN-Spec resin (HDEHP — di-ethylhexil phosphoric acid supported on PTFE powder). Sm and Nd samples were loaded on Re evaporation filaments of double filament assemblies and the isotopic measurements were carried out on a multi-collector Finnigan MAT 262 mass spectrometer in static mode. Uncertainties for Sm/Nd and ¹⁴³Nd/¹⁴⁴Nd ratios are better than ±0.4% (1σ) and ±0.005% (1σ) respectively, based on repeated analyses of international rock standards BHVO-1 and BCR-1. ¹⁴³Nd/¹⁴⁴Nd ratios were normalized to ¹⁴⁶Nd/¹⁴⁴Nd of 0.7219 and the decay constant (λ) used was 6.54 × 10^{–12}.

5.2. U–Pb geochronology of zircons from amphibolite

The dated basic rock consists of orthopyroxene, hornblende, plagioclase, clinopyroxene, biotite, and quartz. Two populations of zircons are distinguished, one comprising colourless, elongate, prismatic, and inclusion free crystals, the other consisting of clean, rounded, red to pink colour crystals.




Fig. 6. U–Pb ages defined by plotting the data in Table 1 and relative to zircons from amphibolite in the largest boudin in the Airport outcrop. Details in text.




Fig. 7. Detail concordia diagram for the three points in the upper part of the diagram in Fig. 6, showing the accurate U–Pb age obtained for metamorphism M_3 in the study outcrop. Details in text.

Table 1 contains a summary of the U–Pb data from the analyses of 13 fractions of grains of zircons. The concordia diagram for the analytical data shows considerable complexity in the U–Pb isotopic system, and the scattered plot of grains of zircons indicates large discordance with the lower and upper intercepts of the diagram (Fig. 6).

We interpret the colourless zircons as representative of the igneous protolith. Coincidentally, their alignment provide the older U–Pb age of 2577 ± 110 Ma among other possible regressions. In despite the large error, this age is similar to the most precise U–Pb SHRIMP age of 2580 ± 10 Ma obtained by Oliveira et al. (2004) for Caraíba norite, suggesting contemporary magmatic events. The population of red-pink zircons (interpreted as metamorphic crystals) plots along another discordia and define an age of 2082 ± 4 Ma that we accept as a good result (Fig. 7) for the regional M_3 metamorphism, because it is very similar to crystallization ages of clearly syntectonic intrusions, such as G_3 granites in the southern part of the Caraíba orebody (2051 ± 16 Ma age, U–Pb from monazite; D'el-Rey Silva et al., 1996), and the Itiúba Syenite (2084 ± 9 Ma U–Pb SHRIMP age; Oliveira et al., 2004). Points probably related to Pb loss plot very near to the lower intercept of the diagram, providing ages of 1.0 and 0.8 Ga that currently we speculate to be geologically meaningless.




Fig. 8. Plotting the data from eight samples of amphibolite in Table 2 in the Sm–Nd diagram results in two isochronic ages. According to Frost and Frost (1995) the causes for moving points out of true isochronic lines may be due to: 1 — Re-equilibrium at outcrop scale that would move the points along the vertical axis ($^{143}\text{Sm}/^{144}\text{Nd}$ ratio), then leading to a new isochronic straight line and affecting the original age of the protolith; 2 — Variation in the igneous protolith at the time of formation, a situation that would increase the $^{147}\text{Sm}/^{143}\text{Nd}$ ratio, changing the inclination of the lines without re-equilibration of the system; and 3 — Re-distribution of Sm relative to Nd during high-grade metamorphism, most likely to be the case in the Airport outcrop, as the process would generate parallel straight lines, but would not re-equilibrate the originally igneous data, therefore the ages will rather correspond to isochronic alignments.

5.3. Sm–Nd geochronology in the Airport outcrop

The systematic study of Nd isotopes was applied to ten samples: eight of them collected in different boudins of amphibolite (1A–H; Fig. 4), one sample of the G_2 tonalite (2A), and one sample of the G_3 granite (3A), all listed in Table 2.

The isotopic data of G_2 tonalite [sample 2A: $T_{\text{DM}} = 3101$ Ma; $\epsilon_{\text{Nd}}(t) = -6.79$] and G_3 granite [sample 3A: $T_{\text{DM}} = 2719$ Ma; $\epsilon_{\text{Nd}}(t) = -9.87$] suggest derivation from partial melting of Mesoarchean continental crust, likely to be the continental part of the basement of the Tanque Novo supracrustals. Plotted together in a single $^{143}\text{Nd}/^{144}\text{Nd}$ versus $^{147}\text{Sm}/^{144}\text{Nd}$ isochronic diagram (Fig. 8), the data from the eight samples of amphibolites, appear to define two isochrones. The older age of 2603 ± 110 Ma, with $\epsilon_{\text{Nd}}(t)$ values close to zero, is interpreted as the age of crystallization of the igneous protolith of the amphibolites. The younger age of 2022 ± 290 Ma and the negative $\epsilon_{\text{Nd}}(t)$ values are consistent with late isotopic re-

Table 2
Summary of the Sm–Nd data from the eight samples collected in the airport outcrop

Sample	Rock Type	Sm (ppm)	Nd (ppm)	$^{147}\text{Sm}/^{144}\text{Nd}$	$^{143}\text{Nd}/^{144}\text{Nd} \pm 2\sigma$	$\epsilon_{(0)}$	T_{DM} (My)	$\epsilon_{\text{Nd}}(t)$
1A	Amphibolite	1.94	7.08	0.1658	0.512124 (12)	−10.02	2961	+0.14
1B	Amphibolite	3.44	13.99	0.1487	0.511724 (09)	−17.83	3126	−2.03
1C	Amphibolite	2.82	8.76	0.1960	0.512619 (24)	−0.37	2884	−0.14
1D	Amphibolite	4.64	19.74	0.1393	0.511648 (17)	−19.31	2869	−0.42
1E	Amphibolite	5.10	20.19	0.1534	0.511816 (25)	−16.04	3104	−1.79
1F	Amphibolite	3.98	16.44	0.1466	0.511855 (32)	−15.28	2705	+1.21
1G	Amphibolite	4.59	17.37	0.1606	0.511930 (22)	−13.81	2918	−1.93
1H	Amphibolite	2.06	7.73	0.1616	0.512015 (23)	−12.15	3048	−0.60
2A	Tonalite (G_2)	7.24	35.39	0.1238	0.511241 (25)	−27.24	3101	−6.79
3A	Granite (G_3)	1.79	19.84	0.0548	0.510202 (19)	−47.52	2719	−9.87




Fig. 9. $\epsilon_{Nd}(t) \times Nd$ content Cartesian graph constructed with data from samples 1A–H in Table 2.

equilibration of the Nd isotopic system and partial re-equilibration of Nd isotopic system during regional metamorphism. This process could have been theoretically related to contaminating fluids with low Sm/Nd ratio (McCulloch and Black, 1984), or loss of Sm relative to Nd during metamorphism.

The values obtained for the Nd isotopes of individual samples of amphibolites indicate a heterogeneous isotope distribution reflected in varied T_{DM} ages and variable values of $\epsilon_{Nd}(t)$ (Fig. 9), and we found this related to the size of the boudins. Samples with values of $\epsilon_{Nd}(t)$ close to zero, or slightly positive (1A, 1C, 1D, and 1H) were all from large boudins (Fig. 4) and are indicative of preservation of the igneous protolith's isotopic composition with T_{DM} model ages around 2.7–2.8 Ga, whereas T_{DM} model ages around 2.9–3.1 Ga

correspond to samples from the border of a boudin (1B); or from small boudins (1E and 1G) with more fractionated Sm/Nd ratios and the most negative $\epsilon_{Nd}(t)$ values that reflect larger isotopic interaction with the host rocks. Sample 1F (from the largest boudin; Fig. 4) displays the best preserved original composition and the highest positive value of $\epsilon_{Nd}(t)$.

6. Discussion

The isotopic data suggest that the geochemical signature of the amphibolites is compatible with basalts sourced from an initially depleted mantle. These, combined with the strongly similar ages of crystallization for Caraiba norites (Oliveira et al., 2004) and airport amphibolite (this paper) leads to consider that the protolith of the Caraiba orebody could have been either: a — basaltic oceanic crust; b — basalts shed as flows within sediments in the Tanque Novo sequence; or c — a gabbroic sill intruded into the Tanque Novo sequence. All these possibilities agree with the large inventory of data from the area surrounding Caraiba, and from the Curaçá terrane as a whole. This inventory requires that before D_1 deformation the orebody's protolith was a conformable body in the same sub-horizontal position of the Tanque Novo volcanic-sedimentary pile, so that all lithotypes above came to experience the same D_1 – D_3 structural/metamorphic evolution.

The age of M_1 metamorphism has not been determined. However, once the field relationships in the Airport outcrop indicate a progressive evolution similar to the one observed in the Caraiba orebody, and the observation that the G_2 intruded syntectonically, all support the interpretation that the structural and metamorphic evolution of the Curaçá terrane may be bracketed between (2.35?)2.3–2.25 and 2.05 Ga (D'el-Rey Silva et al., 1996), with cooling lasting until 1900 Ma ago. The new ages described in this paper suggest a more detailed scenario: M_2 granulite facies conditions peaked in the 2250–2200 Ma interval, whereas M_3 amphibolite facies conditions peaked at 2080–2050 Ma interval (Fig. 10).




Fig. 10. Summary diagrams for the tectonic evolution of the Curaçá belt. D_1 – D_3 structures are observed since cm- to dm-scale up to the scale indicated below each diagram. The leftmost diagram displays G_1 conformable bodies in the Tanque Novo Complex, all affected by S_1 and intrafolial F_1 folds. M_1 metamorphism probably peaked in the 2.3–2.8 Gy interval. The 2.58 Gy old body (Oliveira et al., 2004) destined to be the Caraiba orebody is depicted in the center, with no implication on the origin of the protolith. The central diagram displays S_2 and asymmetric F_2 folds affecting previous units and structures, as well as syntectonic G_2 intrusions, all associated to M_2 metamorphism ($T \leq 1000$ °C; Leite, 2002) peaked in the 2.25–2.20 Gy interval (this paper). The diagram to the right shows D_1 – D_2 units and structures affected by F_3 folds and S_3 , and intruded by G_3 bodies, all affected by M_3 metamorphism peaked 2.08–2.05 Gy ago (our result combined with data in D'el-Rey Silva et al., 1996; Oliveira et al., 2004). Final cooling lasted until ~ 1.9 Gy ago. The Caraiba mushroom is to the left of the Curaçá antiform (Ca). Temperatures around 600 °C for M_1 and M_3 fit data in Leite (2002) for amphibolite facies metamorphism in his PhD study area.

Three similarly important facts make the Curaçá belt a classical area for tectonic analysis based on the combination of accurate field geology, detailed structural observations, and careful isotope analysis of different systems.

Firstly, the large errors associated with the ages in here indicate open-system behaviour of the U–Pb and Sm–Nd isotopic systems in the amphibolite protolith. The difficulty of dating basic rocks in a high grade terrane resides in that such rocks contain extremely small volumes of minerals suitable for high precision age determinations, and their isotopic systems are commonly affected at varied intensities during severe metamorphism, so the study of these systems will indicate varied levels of complexity, no matter if the tectonic evolution was polyphase or polycyclic (Frost and Frost, 1995; Gruau et al., 1996; Pidgeon and Wilde, 1998; Carson et al., 2002).

Secondly, the older cratonic blocks in the São Francisco Craton preserve different isotopic ratios (Fig. 1a) acquired before the Paleoproterozoic. The I–S–C orogen may involve multiple collisions along the margins of the Gavião and Serrinha Mesoarchean blocks during the Neoproterozoic and Paleoproterozoic collisions, and unravelling these presents a considerable challenge.

In the area south of the Itiúba syenite, Teixeira (1997) mapped the Caraíba and Tanque Novo (or Tanque Novo-Ipirá) complexes folded/tectonically imbricate together with the São José do Jacuípe Suite, a unit consisting of norites, gabbro-norites, gabbros, peridotites, and pyroxenites, and suggested that part of the Caraíba Complex defines a magmatic arc, subsequently dated as Neoproterozoic (quoted in Delgado et al., 2003), whereas part of the two other units matches the characteristics of oceanic crust. The interpretation here that the Airport outcrop may preserve 2.6 Ga oceanic crust fits this model well.

The third fact is that the long-lived and transpression-related tectonics recorded in the I–S–C orogen has a great potential to transform a single suture into a stack of sub-vertical suture segments, and the traces of these segments may spread over a wider field area than the original suture zone. In case of polycyclic evolution, this tectonic scenario could have stacked segments of two collision sutures, creating a true geological puzzle.

We are aware that other regressions are possible with our U–Pb data. However, the U–Pb age of 2.6 Gy for the amphibolite protolith is sustained here because: a — it is the highest value obtained; b — it derives from the population of clean zircons; c — it is within error of the Sm–Nd isochronous age we obtained for the same amphibolites; and d — both ages above are similar to the SHRIMP age obtained for the Caraíba norite (Oliveira et al., 2004). Metamorphic zircons in the amphibolites yield an age between 2.3 to 2.2 Ga. We suggest that the 2.3–2.2 Ga interval is the time in which the isotopic system in the amphibolites was disturbed by incoming fluids derived from the G_2 tonalite, and sustain the interpretation based on two facts: 1 — the G_2 tonalite is a syn-tectonic intrusion; and, 2 — pristine zircons from the G_2 tonalite in the same Airport outcrop provided an age of 2248 ± 36 Ma (D'el-Rey Silva et al., 1996), emphasizing the significance of this early Paleoproterozoic D_2 event.

Combining our results with those of previous workers, we propose the following scenario for the tectonic evolution of the region. The protoliths to the Airport amphibolites and the Tanque Novo sequence, both formed ~ 2.6 Ga, when a wide ocean separated the Gavião and Serrinha blocks. The Caraíba Complex was evolving in this ocean as an island arc above an east-dipping subduction zone that plunged below the Serrinha Block (Leite, 2002). M_1 amphibolite facies metamorphism (our results) peaked sometime between 2.35–2.80 Ga, as the rocks entered in the subduction zone, then underwent layer-parallel shearing, and acquired D_1 structures. D_2 Structures developed as the rocks were buried further, and the E–W fold axes (F_2) formed by differential flow affecting fold axes originally trending N–S, a kind of embayment process pointing-down to the east, inside the N–S trending subduction zone and coeval with granulite facies metamorphism (M_2). This metamorphism condition peaked at 2.25–2.2 Ga, our result that on one hand fits in the (2.2–2.1 Ga) interval suggested by Leite (2002) for amalgamation of the island arc to the Serrinha Block, and on the other hand, also suggests a back-arc scenario for part of the Tanque Novo sequence as well as for the amphibolites and the Cu-mineralized orebodies (see also Bello, 1986). Closure of the ocean at ~ 2.1 Ga resulted in the formation of a foreland basin to the west (all in Leite, 2002). The roots of the orogen started to be uplifted as the subduction zone was choked when the Gavião Block began to enter into the subduction zone, and the ultimate Gavião – Serrinha continental collision at 2.08 Ga (Leite, 2002) caused D_3 deformation under M_3 amphibolite facies metamorphic conditions peaked at 2.08–2.5 Ga throughout the I–S–C orogen.

7. Conclusions

This paper has brought to light new U–Pb and Sm–Nd geochronology data obtained after detailed studies carried out in amphibolites (mainly), G_2 tonalite, and G_3 granite of the Airport outcrop, situated ~ 5 km to the north of the Caraíba Cu-orebody, in the Curaçá terrane, northern São Francisco Craton, Brazil. The new data, coupled with well-known tectonic structures and other field data from the Airport outcrop, clearly imply D_1 – D_3 progressive deformational events under amphibolite-granulite-amphibolite M_1 – M_3 metamorphism, and G_1 – G_3 magmatism. The G_2 tonalite and G_3 granite in the Airport outcrop derive (Sm–Nd data) from partial melting of a Mesoarchean continental crust existing in the Curaçá terrane. This Mesoarchean crust is likely to be part of the basement of the Tanque Novo supracrustals. The igneous protolith of the amphibolites in the outcrop is Neoproterozoic and crystallized ca. 2600 Ma ago (U–Pb and Sm–Nd data). This may be also in the age-interval of sedimentation in the northern part of the basin precursor of the Itabuna-Salvador-Curaçá orogen. Granulite facies metamorphism (M_2) and amphibolite facies metamorphism (M_3) in the Curaçá terrane are respectively dated around 2250–2200 Ma, and 2084–2050 Ma (U–Pb data). The progressive evolution supports previous interpretation that M_1 amphibolite facies metamorphism (still lacking a direct age determination) is as old as 2300 Ma.

However, the progressive evolution recorded in the Curaçá terrane as a whole does not rule out the possibility that other parts of the Curaçá terrane, or elsewhere in the I–S–C orogen, still hide evidence for a polycyclic evolution. The results in here emphasize the importance of combining geochronology, lithostructural and metamorphic detailed studies of key outcrops for unravelling the evolution of high-grade terrains, and constitute a contribution for the geochronology approach of high-grade terranes in general.

Acknowledgements

Dr. Sérgio L. Junges is thanked for support during isotopic analyses at the Geochronology Lab — UnB. Elton L. Dantas thanks CNPq for grant 471144/03-7. The first author dedicates his part in this work to his wife and sons, and to the memory of geologist José Genário de Oliveira (deceased in 2000), an enthusiastic of the Caraíba Project, who largely facilitated the visit to the mine and collection of the samples in the Airport outcrop. Two anonymous reviewers are thanked for excellent comments that helped to make better this piece of work. Dr. Alan Collins (The University of Adelaide, Australia) is deeply acknowledged for his superb help on final editing. This paper is a contribution to IGCP project 509, Palaeoproterozoic Supercontinents and Global Evolution.

References

- Ackermand, D., Herd, R.K., Reinhardt, M.C., Windley, B.F., 1987. Sapphirine parageneses from the Caraíba Complex, Bahia, Brazil: the influence of Fe^{2+} – Fe^{3+} distribution on the stability of Sapphirine in natural assemblages. *Journal of Metamorphic Geology* 5, 323–339.
- Barbosa, J.S.F., 1996. O embasamento Arqueano e Proterozóico Inferior de Estado da Bahia. In: Barbosa, J.S.F., Dominguez, J.M.L. (Eds.), *Geologia da Bahia, Texto Explicativo SME-SGM, Capítulo III*, pp. 63–83.
- Barbosa, J.S.F., Sabaté, P., Dominguez, J.M.L., 1996. O estado da Bahia na Plataforma Sul-Americana, suas sub-divisões, critérios de empilhamento estratigráfico das coberturas plataformais e ciclos geotectônicos. In: Barbosa, J.S.F., Dominguez, J.M.L. (Eds.), *Geologia da Bahia, Texto Explicativo SME-SGM, Capítulo II*, pp. 41–61.
- Barbosa, J.S.F., Sabaté, P., 2002. Geological features and the Paleoproterozoic collision of four Archean crustal segments of the São Francisco Craton, Bahia, Brazil. *Anais da Academia Brasileira de Ciências* 74 (2), 343–359.
- Barbosa, J.S.F., Sabaté, P., 2004. Archean and Paleoproterozoic crust of the São Francisco Craton, Bahia, Brazil: Geodynamic Features. *Precambrian Research* 133, 1–27.
- Bello, R.M.S., 1986. Jazida de Cobre de Surubim, Vale do Curacá, Bahia: Mineralogia, Petrografia e Petrogênese. Ph.D. thesis, University of São Paulo (USP).
- Carson, J.C., Ague, J.J.A., Grove, M., Coath, D.C., Harrison, M.T., 2002. U–Pb isotopic behaviour of zircon during upper-amphibolite facies fluid infiltration in the Napier Complex, east Antarctica. *Earth and Planetary Science Letters* 199, 287–310.
- Condie, K.C., 1989. *Plate Tectonics and Crustal Evolution*, Third ed. Pergamon Press, Oxford-UK.
- D'el-Rey Silva, L.J.H., 1984. Geologia e controle estrutural do depósito cuprífero Caraíba, Vale do Curaçá, Bahia, Brasil. M.Sc. thesis, Federal University of Bahia (UFBA).
- D'el-Rey Silva, L.J.H., 1985. Geologia e controle estrutural do depósito cuprífero Caraíba, Vale do Curaçá, Bahia. *Geologia e Recursos Naturais do Estado da Bahia, SME, Série Textos Básicos* 6, 51–123.
- D'el-Rey Silva, L.J.H., 1993. Geologia Estrutural e Tectônica na área do Projeto Lajedinho-Ipirá. Consulting report for CBPM, Salvador, 23 pp., 23 figures.
- D'el-Rey Silva, L.J.H., Oliveira, J.G., 1999. Geology of the Caraíba Copper Mine and its surroundings in the Paleoproterozoic Curaçá Belt–Curaçá River Valley, Bahia, Brazil. In: da Silva, M. da G., Misi, A. (Coordenadores) *Base metal Deposits of Brazil, MME/CPRM/DNPM*, 25–32.
- D'el-Rey Silva, L.J.H., Cavalcante, P.R.B., Mota, E.R., Rocha da, A.M.R., 1988. Controle estrutural da mina de cobre Caraiba: implicações na lavra e na tectônica das faixas móveis do Proterozóico Inferior. XXXV Congresso Brasileiro de Geologia, Belém, Brazil, *Anais SBG*, vol. 1, pp. 16–29.
- D'el-Rey Silva, L.J.H., Oliveira, J.G., Lima e Silva, F.J., 1994. The Mushroom-shaped Caraíba Cu-deposit, Vale do Curaçá-Ba: Understanding the Structural Evolution of the Paleoproterozoic, Granulitic Ipirá-Curaçá Belt within the São Francisco Craton. XXXVIII Congresso Brasileiro de Geologia, Camboriú, Brazil, *SBG, Boletim de Resumos Expandidos*, vol. 2, pp. 175–177.
- D'el-Rey Silva, L.J.H., Oliveira, J.G., Gaál, E.G., 1996. Implication of the Caraíba Deposit's structural controls on the emplacement of the Cu-bearing hyperthenites of the Curaçá Valley, Bahia-Brazil. *Revista Brasileira de Geociências* 26 (3), 181–196.
- Delgado, I. de M., Souza, J.D., 1975. Projeto cobre Curaçá. *Geologia Econômica do Distrito Cuprífero do Rio Curaçá — Bahia, Brasil. CPRM/DNPM*, internal report, 30 volumes.
- Delgado, I. de M., de Souza, J.D., da Silva, L.C., Silveira Filho, N.C., dos Santos, R.A., Pedreira, A.J., Guimarães, J.T., Angelim, L.A. de A., Vasconcelos, A.M., Gomes, I.P., de Lacerda Filho, J.V., Valente, C.R., Perrotta, M.M., Heineck, C.A., 2003. *Geotectonics of the Atlantic Shield. Geology, Tectonics and Mineral Resources of Brazil: Text, Maps, and GIS, CPRM*, pp. 227–334. 692 p., Chapter V.
- Figueiredo, M.C.H., 1981. Geoquímica da rochas metamórficas de alto grau do nordeste da Bahia-Brazil. *Geologia e Recursos Naturais do Estado da Bahia, SME, Série Textos Básicos* 4, 1–71.
- Frost, C.D., Frost, B.R., 1995. Open-system dehydration of amphibolites, Morton Pass, Wyoming: elemental and Nd and Sr isotopic effects. *The Journal of Geology* 103, 269–284.
- Gaal, E.G., 1982. Evaluation of geological, geophysical, and geochemical data of the Curaçá Valley, and the Mundo Novo-Uauá-Rio Capim area. Consulting report for the Mineração Caraíba Ltda, MID-066/82.
- Gioia, S.M.L., Pimentel, M.M., 2000. The Sm–Nd method in the geochronology laboratory of the University of Brasília. *Anais da Academia Brasileira de Ciências* 72, 219–245.
- Gromet, L.D., 1991. Direct dating of deformational fabrics. In: Heaman, L., Ludden, J.N. (Eds.), *Short Course Handbook on Applications of Radiogenic Isotope Systems to Problems in Geology*. Mineralogists Association of Canada, pp. 167–192. Chapter 5.
- Gruau, G., Rosing, M., Bridgwater, D., Gill, R.C.O., 1996. Resetting of Sm–Nd systematics during metamorphism of >3.7 Gy rocks: implications for isotopic models of early Earth differentiation. *Chemical Geology* 133, 225–240.
- Hasuy, Y., D'el-Rey Silva, L.J.H., Lima e Silva, F.J., Mandetta, P., de Moraes, J.A., Oliveira, J.G., Miola, W., 1982. Geology and copper mineralization of Curaçá River Valley, Bahia. *Revista Brasileira de Geociências* 12 (1-3), 463–474.
- Inda, H.A.V., Barbosa, J.S.F., 1978. Mapa geológico do Estado da Bahia, escala 1:1.000.000. Texto explicativo. SME/CPM.
- Jardim de Sá, E.F., Arcanjo, C.J., Legrand, J.-M., 1982. Structural and metamorphic history of part of the high-grade terrane in the Curaçá Valley, Bahia, Brazil. *Revista Brasileira de Geociências* 12, 251–262.
- Krogh, T.E., 1973. A low-contamination method for hydrothermal decomposition of zircon and extraction of U and Pb for isotopic age determinations. *Geochimica et Cosmochimica Acta* 37, 485–494.
- Leite, C.M.M., 2002. A Evolução Geodinâmica da Orogênese Paleoproterozóica nas Regiões de Capim Grosso-Jacobina e Pintadas-Mundo Novo (Bahia-Brasil): Metamorfismo, Anatexia Crustal e Tectônica. Ph.D. thesis, Federal University of Bahia (UFBA).
- Leite, C.M.M., Barbosa, J.S.F., Nicollet, C., Kienast, J.R., Fuck, R.A., 2005. Ultrahigh-temperature metamorphism of sapphirine-bearing granulite from the 2.0 Gy Itabuna-Salvador-Curaçá orogen, Bahia, Brazil. III Simpósio sobre o Cráton do São Francisco, Salvador, Brasil, *AnaisISSN :1808-6047*, pp. 83–86.
- Lindenmayer, Z.G., 1981. Evolução geológica do Vale do Curaçá e dos corpos máfico-ultramáficos mineralizados a cobre. M.Sc. thesis, Federal University of Bahia (UFBA).

- Lindenmayer, Z.G. and six co-authors, 1984. Projeto Cu Bahia, Alvará 872, DNPM 812.998/73. Relatório Final de Pesquisa, DOCEGEO-CVRD, 3 volumes.
- Ludwig, K.R., 1993. PBDAT. A computer program for processing Pb–U–Th isotope data. USGS Open File Report, vol. 88-542. 34 pp.
- Ludwig, K.R., 2001. Users Manual for Isoplot/Ex version 2.47. A geochronological toolkit for Microsoft Excel. Berkeley Geochronology Centre Special Publication, vol. 1a. 55 pp.
- Mandetta, P., 1982. Aspectos geológicos e petrogenéticos das associações máfico-ultramáficas da região de Caraíba, Vale do Curacá/Ba. MsC thesis, Federal University of Bahia (UFBA).
- Mayer, W.D., Barnes, S.-J., 1996. Unusually high concentrations of magnetite at Caraíba and other Cu-sulphide deposits in the Curaçá Valley, Bahia, Brazil. *The Canadian Mineralogist* 34, 717–731.
- McCulloch, M.T., Black, L.P., 1984. Sm–Nd isotopic systematics of Enderby Land granulites and evidence for the redistribution of Sm and Nd during metamorphism. *Earth and Planetary Science Letters* 71, 46–58.
- Oliveira, E.P., Windley, B.F., McNaughton, N.J., Pimentel, M.M., Fletcher, I.R., 2004. Contrasting copper and chromium metallogenic evolution of terranes in the Paleoproterozoic Itabuna-Salvador-Curaçá orogen, São Francisco craton, Brazil: new zircon (SHRIMP) and Sm–Nd (model) ages and their significance for orogen-parallel escape tectonics. *Precambrian Research* 128, 143–165.
- Passchier, C.W., Myers, J.S., Kröner, A., 1990. *Field Geology of High-Grade Gneiss Terranes*. Springer-Verlag, Berlin.
- Pidgeon, R.T., Wilde, S.A., 1998. The interpretation of complex zircon U–Pb systems in Archean granitoids and gneisses from the Jack Hills, Narryer Gneiss Terrane, Western Australia. *Precambrian Research* 91, 309–322.
- Ramsay, J.G., 1967. *Folding and Fracturing of Rocks*. McGraw Hill, New York.
- Sabaté, P., Marinho, M.M., Vidal, P., Caen-Vachette, M., 1990. The 2-Ga peraluminous magmatism of the Jacobina-Contendas Mirante belts (Bahia, Brazil): geological and isotopic constraints on the sources. *Chemical Geology* 83, 325–338.
- da Silva, M. da G., 1996. Sequências metasedimentares, vulcanosedimentares e greenstone belts do Arqueano e Proterozóico Inferior. In: Barbosa, J.S.F., Dominguez, J.M.L. (Eds.), *Geologia da Bahia, Texto Explicativo SME-SGM, Capítulo IV*, pp. 86–102.
- Teixeira, L., 1997. O complexo Caraíba e a suíte São José do Jacuípe no cinturão Salvador-Curaçá (Bahia, Brasil): petrologia, geoquímica e potencial metalogenético. Ph.D. Thesis, Federal University of Bahia (UFBA).
- Windley, B.F., 1986. *The Evolving Continents*, Second ed. John Wiley and Sons Ltd, UK.