

Seroprevalence and Risk Factors associated with *Helicobacter pylori* Infection in Blood Donors in Salvador, Northeast-Brazil

Andre C. Lyra¹, Genoile Santana¹, Nelma Santana³, Anibal Silvany-Neto², Emilia Magalhães¹, Eduardo M. Pereira⁴, Ramiro Mascarenhas¹, Marcos C. Lyra¹, Andrea Veiga¹, Karina Ferreira¹, Schilioma Zaterka⁵ and Luiz G. Lyra¹

Gastroenterology and Hepatology Service of Federal University of Bahia and Gastro-Hepatology Service of São Rafael Hospital¹, Department of Epidemiology of Federal University of Bahia², Hemocentro of Bahia (HEMOBA)³, Central Laboratory – LACEN⁴, Salvador, BA; Discipline of Gastroenterology⁵, Unicamp, SP, Brazil

Helicobacter pylori plays an important role in the etiology of peptic ulcer disease. Its prevalence appears to be higher in developing countries. We evaluated the seroprevalence of *H. pylori* and risk factors associated with infection in voluntary blood donors who attended the main blood center of the city of Salvador, Brazil. The subjects responded to an epidemiological questionnaire, with information about sex, age, race, lifestyle, social-economic level indicators, and residence and hygiene conditions. Anti-*H. pylori* antibody was determined by ELISA (Cobas Core, Roche). Three hundred and seven subjects were included in the study. Anti-*H. pylori* antibody results were indeterminate in 33 individuals (10.8%), who were excluded from analysis. Among the remaining 274 subjects, 187 (68.2%) were anti-*H. pylori* positive. Based on multivariate logistic regression analysis three variables were found to be significantly associated with a higher prevalence of *H. pylori* infection: absence of plumbing in the residence during childhood, a history of rainwater invading the dwelling during childhood, and low ingestion of milk.

Key Words: *H. pylori*, seroprevalence, risk factors, Brazil.

Helicobacter pylori plays a significant role in the etiology and pathogenesis of chronic active gastritis and peptic ulcer disease [1,2,3]. It is estimated that 50% of the world's population is infected by this bacteria [4,5]. The microorganism was first described in 1983 [6,7], but its epidemiology has not been entirely elucidated. Data regarding risk factors and modes of transmission are still controversial; it appears that there could be several modes of transmission [8,9,10]. In Brazil, peptic

ulcer disease is an important cause of morbidity [11]. There are several studies about the epidemiology of *H. pylori* infection in the Brazilian population. Rocha et al. [12], using indirect immunofluorescence, detected a prevalence of 62.1% *H. pylori* infection in asymptomatic Brazilian blood donors in an urban area, this is a less specific test than the enzyme-linked immunosorbent assay (ELISA). Souto et al. [13] reported a prevalence of 84.7% *H. pylori* infection in adults in a rural area in a central region of Brazil, using ELISA. Further studies regarding the epidemiology of this bacterial infection in the Brazilian population should be encouraged in order to identify local risk factors. We evaluated the prevalence of *H. pylori* infection in voluntary blood donors of the city of Salvador in Northeast Brazil, utilizing an enzyme-linked immunosorbent assay (ELISA- COBAS CORE anti-*H.pylori*), and we identified possible risk factors.

Received on 20 August 2002; revised 10 October 2003.

Address for correspondence: Dr. Andre C Lyra, MD, Serviço de Gastro-Hepatologia do Hospital São Rafael, Salvador-BA, Brasil. Rua Socrates Guanaes Gomes, 84, room 401, Salvador – Bahia – Brazil – Zip code: 40.283.320 Phone – Fax (55)-(71)-452-9589. This study was supported in part by the CNPq.

The Brazilian Journal of Infectious Diseases 2003;7(5):339-345
© 2003 by The Brazilian Journal of Infectious Diseases and Contexto Publishing. All rights reserved.

Materials and Methods

Subjects. We made a cross-sectional study of voluntary blood donors who attended the main blood center of Bahia (HEMOBA) and who accepted participation in the study during 1997 and 1999. Informed written consent was obtained from all subjects; the study was approved by the institutional review boards and the protocol conformed to the ethical guidelines of the 1975 Declaration of Helsinki.

Inclusion and exclusion criteria. We only included patients who were born and who lived in the city of Salvador, Bahia; they must have fulfilled the blood center criteria for blood donation, which included, among other requirements, being 18 to 60-years-old. Patients who had lived in any city other than Salvador for any period of time were excluded from the study.

Variables analyzed. Subjects responded to an epidemiological questionnaire immediately before blood donation. The questionnaires were applied on weekdays in a random fashion. No apparent unusual situation in the blood center was detected during the period the interviews were done. The independent variables studied were: gender, race, weight, height, age (over or under 28), civil state (married / living together, or single), alcohol and coffee consumption, smoking, type and frequency of food ingested (beef, lamb, poultry, fish, pork, vegetables, fruits, milk), type of water ingested during childhood and adulthood (filtered / boiled, or not), garbage collection in the neighborhood during childhood and adulthood, indicators of social-economic status: educational level (complete high school or higher level; incomplete high school or lower level), family income (over or less than 5 minimum salaries), neighborhood (wealthy or not), type of dwelling (shack or house / apartment), water supply (presence or absence of plumbing), sewage system, number of rooms and number of persons residing in each dwelling, history of rainwater invading or not the dwelling (the latter four items were asked regarding childhood and present time). No main independent variable was specified due to the

exploratory nature of the study. *Helicobacter pylori* infection was considered the dependent variable.

ELISA for *H. pylori*. At the time of venous puncture for donation a 10-ml blood sample was drawn and the sera were stored at -20°C . The ELISA for *H. pylori* was performed after all questionnaires were given. The period between the time the first serum was stored and the ELISA tests was 14 months. Anti-*H. pylori* antibodies were assayed by the Cobas Core anti-*H. pylori* IgG EIA (Roche Diagnostics Division, Basel-Switzerland), a second generation EIA. The test was performed essentially according to manufacturer's instructions. This method has been previously validated in the Brazilian population. The sensitivity, specificity and positive and negative predictive values of the test are 95.4, 100, 100 and 91.4%, respectively [14]. Indeterminate results were excluded from analysis.

Statistical analysis. The sample size was obtained as previously described [15]. We considered an N of 45,000 (estimated number of donors in one year), an absolute precision of 5.5%, a confidence level of 95%, and an estimated prevalence of *H. pylori* infection of 68.2%. This prevalence was obtained retrospectively at the end of the study. Using these parameters, the required sample size should be 274 donors. The chi-square test was used to compare discrete variables; continuous variables were expressed as mean \pm SEM and were analyzed by a non-paired two-tailed t-test with a significance level of 5%. Levenes' test was used to assess variance homogeneity. A multiple logistic regression analysis was carried out in an exploratory sense in order to adequately identify important independent variables. Prevalence ratios of infection were used as measures of association during the simple analysis, and odds ratios of infection in the logistic regression analysis, given the elevated frequency of the *H. pylori* infection in the population. Interaction terms were not taken into account. Variables were selected using the backward stepwise method and the statistical inference was based on likelihood ratio tests. The criterium utilized for variable exclusion was a P value

greater than or equal to 0.17. All the categorical independent variables were coded using an indicator variable scheme. The regression considered comparisons of predicted with observed outcomes, the likelihood of the observed results, the goodness-of-fit statistic and a residual analysis. Confidence intervals for odds ratios were calculated. All these statistics were done using the SPSS software (SPSS version 9.0, SPSS Inc. Chicago, IL).

Results

Three hundred and seven blood donors were included in the study. Detection of anti-*H. pylori* antibody was indeterminate in 33 subjects (10.8%). These cases were excluded from analysis. Comparison between this group and the remaining 274 cases showed significant differences for three variables (Table 1). Among the remaining 274 blood donors, 54 (19.7%) were female and 220 (80.3%) were male. The mean age was 29 ± 8.4 years. Fifty one percent were single. Only 112 donors (40.8%) had completed high school. Sixty three percent had an income between one and five minimum salaries, and 89% received up to 12 minimum salaries. The mean number of people residing in a dwelling during childhood was 7.5. The prevalence of anti-*H. pylori* in this population was 68.2%. The prevalence of the antibody in the male group was 71.4% and in the female group it was 55.6% (Table 2). Based on bivariate simple analysis the variables that were significantly associated with the *H. pylori* infection are displayed in Table 3. There was no significant association with race, age, weight, height, civil state, alcohol and coffee consumption, smoking, educational level, family income, garbage collection in the neighborhood during childhood and adulthood, neighborhood, number of rooms and number of persons residing in each dwelling during childhood or at the present time, or water supply and sewage system at the present time.

In the multivariate analysis four variables were significantly associated with *H. pylori* infection at the 5% level: absence of plumbing in the residence during childhood, history of rainwater invading the dwelling

during childhood, low ingestion of milk and high consumption of fish. The variables beef consumption, civil status, current type of dwelling and vegetable consumption showed borderline P values (considering $\alpha = 5\%$) and were not discarded as unimportant (Table 4). Only three variables were significantly associated with a higher prevalence of *H. pylori* infection in both bivariate and multivariate analysis: absence of plumbing in the residence during childhood, history of rainwater invading the dwelling during childhood, and low ingestion of milk.

Discussion

In studying the prevalence of *H. pylori* infection we found that 68% of the subjects were *H. pylori* seropositive by ELISA. The study population consisted of blood donors and thus it does not necessarily correspond to the entire population of the city of Salvador, Bahia. However, the results are in accordance with previous studies that show a high seroprevalence of *H. pylori* infection in Brazilian populations [12,13].

We did not find an association between *H. pylori* infection and educational level, family income or age. Nevertheless, it is important to note that most of the subjects who were enrolled had a low educational level and a low income, and this might have influenced the results. The population also mostly consisted of young adults, with few donors being 50 years-old or over, and 80% of the subjects were male; therefore, this study may not be ideal for an analysis of the whole population. Zaterka et al. found that 70% of the children of low-income families in Brazil are already infected with *H. pylori* when they are 5 years old [16].

Three variables appeared to be risk factors for *H. pylori* infection in both bivariate simple analysis and multivariate analysis: history of rainwater invading the dwelling during childhood, absence of plumbing water system at the residence during childhood, and low ingestion of milk. To our knowledge, this is the first study to suggest rainwater infiltration as a potential source of *H. pylori* infection. This seems to be a plausible source of infection in developing countries,

Table 1. Comparison between the group of 33 blood donors excluded from analysis and the remaining 274 blood donors

Variable	Included cases	Excluded cases N (%)	X ² N (%)	P value
Educational level				
Complete high school or higher educational level	112 (40.9)	24 (72.7)	10.8	0.001
Incomplete high school or lower educational level	162 (59.1)	9 (27.3)		
Garbage collection in the neighborhood at childhood				
Yes	135 (52.7)	25 (75.8)	5.4	0.02
No	121 (47.3)	8 (24.2)		
Type of water ingested at adulthood				
Filtered or boiled	187 (68.2)	31 (93.9)	8.2	0.004
Not filtered or boiled	87 (31.8)	2 (6.1)		

Table 2. Prevalence of anti-*H. Pylori* antibody in 274 blood donors according to gender

Gender	Anti- <i>H. pylori</i>		Prevalence ratio	95% Confidence interval
	Positive	Negative		
Male	157 (71.4%)	63 (28.6%)	1.28	(1.00; 1.65)
Female	30 (55.6%)	24 (44.4%)		
Total	187 (68.2%)	87 (31.8%)		

X² = 4.3 P = 0.038. PR = Prevalence ratio.

such as Brazil, since some neighborhoods may be subject to flooding during heavy rain and this favors the contact between the residents and the material originated from the surrounding sewer system. The association with absence of plumbing in the residence also suggests that *H. pylori* contamination and transmission could occur through water. Water has been considered a possible transmission route of *H. pylori* infection, especially in developing countries [17]. Swimming in rivers, streams, or pools increases the odds of infection, as does using streams as a drinking water source.

In a population-based study that evaluated children from Peru, water supply was an important source of

H. pylori infection in families with high or low social economic levels [18]. The fact the these conditions occurred during childhood is in accordance with previous studies that showed a greater probability of acquiring the infection during childhood; this could be a result of hygiene habits and a higher susceptibility to *H. pylori* infection during this period of life [19,20].

In contrast to a previous study that did not find any association between high or low ingestion of milk and *H. pylori* infection, we found that low milk ingestion was a risk factor for *H. pylori* infection [21]. We also found that alcohol consumption and smoking were not important risk factors; this is in agreement with previous studies. Nevertheless this issue remains controversial [22-25].

Table 3. Distribution of dichotomized variables that were significant based on bivariate analysis (n=274)

Risk factor	N	Anti- <i>H. pylori</i> antibody		Prevalence ratio 95%	Confidence interval
		Positive	Negative		
Rainwater invading dwelling in childhood	270				
Yes		56 (83.6)	11 (16.4)	1.32	(1.13; 1.53)
No		129 (63.5)	74 (36.5)		
Poultry consumption	274				
Yes		46 (55.4)	37 (44.6)	1.33	(1.08; 1.64)
No		141 (73.8)	50 (26.2)		
Presence of plumbing water system in childhood	266				
No		31 (83.8)	6 (16.2)	1.28	(1.08; 1.52)
Yes		150 (65.5)	79 (35.5)		
Sewage system at childhood	263				
No		47 (82.5)	10 (17.5)	1.27	(1.08; 1.48)
Yes		134 (65.0)	72 (35.0)		
Type of water ingested at adulthood	274				
Not filtered nor boiled		68 (78.2)	19 (21.8)	1.23	(1.05; 1.43)
Filtered or boiled		119 (63.9)	68 (36.4)		
Fruit consumption	274				
At least once a week		142 (65.1)	76 (34.9)	0.81	(0.69; 0.95)
Less than once a week		45 (80.4)	11 (19.6)		
Milk ingestion	272				
At least once a week		157 (65.7)	82 (34.3)	0.74	(0.64; 0.87)
Less than once a week		29 (87.9)	4 (12.1)		
Vegetable consumption	274				
At least once a week		132 (63.8)	75 (36.2)	0.78	(0.67; 0.90)
Less than once a week		55 (82.1)	12 (17.9)		

Finally, we initially included 307 patients in the study, however 33 cases had indeterminate *H. pylori* serology results, and thus were excluded from analysis. Although we have found significant differences in three variables between this group and the remaining 274 cases, we believe that the number of excluded cases is probably too low to have influenced the results. Moreover, only two of these variables were previously reported to be associated with *H. pylori* infection. Most probably, the only

possible consequences are an underestimate of prevalence ratios of these variables.

In summary, this study evaluated the seroprevalence of *H. pylori* in a population of blood donors from the city of Salvador in Northeast-Brazil and suggests possible risk factors for *H. pylori* infection in a developing country. Further large scale population-based studies are needed to better determine the role of these potential sources of transmission of *H. pylori*.

Table 4. Risk factors for *H. pylori* infection: results of the multivariate analysis

Covariable	β	SE	Likelihood ratio Test-P-value	Odds Ratio	CI (95%)
Rainwater invading dwelling in childhood					
Yes vs. no	0.9989	0.3986	0.0081	2.7152	(1.24; 5.95)
Beef consumption					
Yes vs. no	0.5700	0.3377	0.0919	1.7682	(0.91; 3.12)
Plumbing water system in childhood					
No vs. yes	1.0661	0.5097	0.0251	2.9040	(1.06; 7.88)
Civil state					
Other vs single	-0.5543	0.3039	0.0663	0.5745	(0.31; 1.04)
Milk ingestion					
At least once a week vs. less than once a week	-1.8832	0.7625	0.0023	0.1521	(0.03; 0.67)
Fish consumption					
Yes vs. no	0.8630	0.3063	0.0042	2.3703	(1.30; 4.32)
Current type of dwelling					
House vs. apartment	0.6426	0.3714	0.0844	1.9015	(0.91; 3.93)
Vegetable Consumption					
Daily or weekly consumption vs. no or biweekly consumption consumption consumption	-0.6970	0.3926	0.0667	0.4981	(0.23; 1.07)
Constant	1.6734	0.9172	0.0681 ^a		

a – Wald's statistic; CI: confidence interval.

Acknowledgements

This study was supported in part by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq).

References

1. Tytgat G.N.J. Treatment of peptic ulcer. *Digestion* **1993**;59:446-52.
2. Moss S.F., Fendrick A.M., Cave D.R., Modlin I.M. *Helicobacter pylori* – More light, less heat. *Am J Gastroenterol* **1998**;93:306-10.
3. Peek Jr. R.M., Blaser M.J. Pathophysiology of *Helicobacter pylori*-induced gastritis and peptic ulcer disease. *Am J Med* **1997**;102:200-7.
4. Megraud F., Brassens-Rabbe M.P., Denis F. Seroepidemiology of *Campylobacter pylori* infection in various populations. *J Clin Microbiol* **1989**;27:1870-3.
5. Perez-Perez G.I., Dwarkin B.M., Chados J.E., Blaser M.J. *Campylobacter pylori* antibodies in humans. *Ann Int Medicine* **1988**;109:11-7.
6. Warren J.R. Unidentified curved bacilli in the stomach of patients with gastritis and peptic ulceration. *Lancet* **1983**;1:1273.
7. Marshall B.J. Unidentified curved bacilli in the stomach of patients with gastritis and peptic ulceration. *Lancet* **1983**;1:1273-4.
8. Hopkins R.J., Vial P.A., Ferreccio C., et al. Seroprevalence of *Helicobacter pylori* in Chile: Vegetables may serve as one route of transmission. *J Infect Dis* **1993**;168:222-6.
9. Mégraud F. Transmission of *Helicobacter pylori*: Faecal-oral versus oral-oral route. *Aliment Pharmacol Ther* **1995**;9(suppl.2):85-91.

10. Rouland M. Transmission of *Helicobacter pylori*: Is it all child's play? *Lancet* **2000**;55:332-3.
11. Oliveira L.C.M., Buso A.G., Siqueira F.º L., et al. Peptic disease and *Helicobacter pylori* are highly prevalent in patients with the indeterminate form of Chagas disease: Report of 21 cases. *Revista do Instituto de Medicina Tropical de São Paulo* **1997**;39:209-12.
12. Rocha G.A., Queiroz D.M.M., Mendes E.N., et al. Indirect immunofluorescence determination of the frequency of anti-*H. pylori* antibodies in Brazilian blood donors *Braz J Med Biol Res* **1992**;25:683-9.
13. Souto F.J.D., Fontes C.J.F., Rocha G.A., et al. Prevalence of *Helicobacter pylori* infection in a rural area of the State of Mato Grosso, Brazil. *Mem Inst Oswaldo Cruz* **1998**;93:171-4.
14. Rocha G.A., Oliveira A.M., Queiroz D.M., et al. Serodiagnosis of *Helicobacter pylori* infection by Cobas Core ELISA in adults from Minas Gerais, Brazil. *Braz J Med Biol Res* **1998**;31:1263-8.
15. Daniel W.W. *Biostatistics: A foundation for analysis in the health sciences*, **1999**; 7th ed – John Wiley & Sons, New York.
16. Zaterka S., Nishimura N.F., Parente J.M., Libanês L. Prevalence of *Helicobacter pylori* infection in children population in Brazil. *Gut* **2002**;51(Suppl III): A93.
17. Suerbaum S., Michetti P. *Helicobacter pylori* infection. *N Engl J Med* **2002**;347:1175-86.
18. Klein P.D., Graham D.Y., Gaillour A., et al. Water source as risk factor for *Helicobacter pylori* infection in Peruvian children. *Lancet* **1991**;337:1503-6.
19. Neale K.R., Logan R.P.H. The epidemiology and transmission of *Helicobacter pylori* infection in children. *Aliment Pharmacol Ther* **1995**;9 (suppl.2):77-84.
20. Oliveira A.M.R., Queiroz D.M.M., Rocha G.A., Mendes E.N. Seroprevalence of *Helicobacter pylori* infection in children of low socioeconomic level in Belo Horizonte, Brazil. *Am J Gastroenterol* **1994**;89:2201-4.
21. Begue R.E., Gonzalez J.L., Correa-Garcian H., Tang S.C. Dietary risk factors associated with the transmission of *Helicobacter pylori* in Lima, Peru. *Am J Trop Med Hyg* **1998**;59:637-40.
22. Ogihara A., Kikuchi S., Hasegawa A., et al. Relationship between *Helicobacter pylori* infection and smoking and drinking habits. *J Gastroenterol Hepatol* **2000**;15:271-6.
23. Everhart J.E., Kruszon-Moran D., Perez-Perez G.I., et al. Seroprevalence and ethnic differences in *Helicobacter pylori* infection among adults in the United States. *J Infect Dis* **2000**;181:1359-63.
24. Woodward M., Morrison C., McColl K. An investigation into factors associated with *Helicobacter pylori* infection. *J Clin Epidemiol* **2000**;53:175-81.
25. Giannuzzi F., Giannuzzi U., Bianciardi L., et al. Risk factors for acquiring *Helicobacter pylori* infection in a group of Tuscan teenagers. *New Microbiol* **2001**;24:165-70.